

A PERTURBATION PROBLEM FOR THE SHIFT SEMIGROUP

MASAKI IZUMI

ABSTRACT. Motivated by B. Tsirelson's construction of E_0 -semigroups of type III, we investigate a C_0 -semigroup acting on the space of square integrable functions of the half line whose difference from the shift semigroup is a Hilbert-Schmidt operator. We give a description of such semigroups in terms of analytic functions on the right-half plane and construct several examples.

1. INTRODUCTION

An E_0 -semigroup is a semigroup of unital endomorphisms of $\mathbb{B}(H)$, the set of bounded operators on an infinite dimensional separable Hilbert space H , with appropriate continuity. Despite the fact that $\mathbb{B}(H)$ is the simplest infinite dimensional factor, the classification of E_0 -semigroups is far from a satisfactory stage mainly caused by the presence of so called type II and type III examples [11],[12],[17]. Indeed, these classes are related to various areas of analysis, such as probability theory and harmonic analysis, which makes the subject more interesting and worth investigating than one would expect at first sight. The reader is referred to Arveson's monograph [3] and contributions in [13], in particular those of Arveson, Powers, and Tsirelson among others, for history and basic results for the subject.

In this paper, we investigate a purely operator theoretical problem motivated by Tsirelson's construction of uncountably many mutually non-cocycle conjugate E_0 -semigroups of type III [18]. Namely, let $\{S_t\}_{t \geq 0}$ be the shift semigroup of $L^2(0, \infty)$, that is

$$(S_t f)(x) = \begin{cases} 0 & (x < t) \\ f(x - t) & (t \leq x) \end{cases}.$$

We investigate the structure of a C_0 -semigroup of bounded operators $\{T_t\}_{t \geq 0}$ acting on $L^2(0, \infty)$ satisfying the following two conditions:

- (C1) $T_t^* S_t = I$ for all $t \geq 0$.
- (C2) $T_t - S_t$ is a Hilbert-Schmidt class operator for all $t \geq 0$.

If T_t as above preserves the real functions $L^2(0, \infty)_{\mathbb{R}}$ in $L^2(0, \infty)$, then a standard argument for quasi-equivalence of representations of the CCR (canonical commutation relation) algebra tells the following: there exists an E_0 -semigroup α_t acting on the symmetric Fock space over $L^2(0, \infty)$ determined by

$$\alpha_t(W(f + ig)) = W(S_t f + i T_t g), \quad f, g \in L^2(0, \infty)_{\mathbb{R}},$$

where $W(f + ig)$ is the Weyl operator for $f + ig$. Although Tsirelson's construction is based on a homogeneous continuous product of measure classes arising from an off

Work supported by JSPS.

white noise [17], his examples actually come from the above construction as we will see in Section 6.

A careful examination of the generator of a C_0 -semigroup $\{T_t\}_{t \geq 0}$ satisfying (C1) shows that $\{T_t\}_{t \geq 0}$ is completely characterized by a holomorphic function $M(z)$ on the right half plane $\mathbb{H}_r$ such that $M(z)/(1+z)$ belongs to the Hardy space $H^2(\mathbb{H}_r)$. We call $M(z)$ the half-density function for $\{T_t\}_{t \geq 0}$. When $\{T_t\}_{t \geq 0}$ comes from an off white noise, the absolute value of the boundary value function $M(iy)$ on the imaginary axis is nothing but the square root of the spectral density function discussed by Tsirelson [17],[18]. We completely characterize C_0 -semigroups satisfying (C1) and (C2) in terms of the function $M(z)$ (Theorem 4.2). It turns out that this class of functions $|M(i\lambda)|^2$ is slightly larger than the class of spectral density functions discussed in [18].

In Section 5, we construct a class of C_0 -semigroups satisfying (C1) and (C2) parameterized by a function φ in the space $L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$. More precisely, we consider the case with $M(z) = 1 - \mathcal{L}[\varphi](z)$ where $\mathcal{L}[\varphi]$ is the Laplace transform of φ . While a perturbation argument shows that the E_0 -semigroups arising from this class of functions are cocycle conjugate to those coming from a subclass of off white noises, one cannot distinguish them from the CCR flow of index 1 by using Tsirelson's invariants discussed in [17] because the spectral density functions $|M(iy)|^2$ converge to 1 at infinity in this case. Yet, in the forthcoming paper [9], we show that this class of E_0 -semigroups contains uncountably many mutually non cocycle conjugate type III E_0 -semigroups. The invariant we adopt for differentiating these examples is the Murray-von Neumann type of the "local observable algebras" for an open subset U of the interval $[0, 1]$, which may be an AFD type III factor if φ does not belong to $L^2(0, \infty)$ and U has a sufficiently complicated shape. It should be noted that the space $L^2((0, \infty), (1 \wedge x)dx)$ also plays a crucial role in Powers' construction of uncountably many mutually non cocycle conjugate E_0 -semigroups of type II_0 [12] though we don't know if there exists a direct relationship between our argument with Powers' in the present stage.

The author would like to thank W. Arveson and R. Srinivasan for stimulating discussions.

2. PRELIMINARIES

Throughout this note, the symbol $\|\cdot\|$ means either the L^2 norm or the operator norm, depending on the context. The symbol $\|\cdot\|_p$ denotes the L^p norm. The Hilbert-Schmidt norm is denoted by $\|\cdot\|_{H.S.}$. Every function space we discuss is complex valued.

For $f, g \in L^2(0, \infty)$, we denote by $\langle f, g \rangle$ the usual inner product

$$\langle f, g \rangle = \int_0^\infty f(x) \overline{g(x)} dx.$$

Dual pairing in various contexts will be denoted by (f, g) . For example, for $f, g \in L^2(0, \infty)$

$$(f, g) = \int_0^\infty f(x) g(x) dx.$$

For an operator A of $L^2(0, \infty)$, we denote by $D(A)$ the domain of A and by $\mathcal{G}(A)$ the graph of A , that is

$$\mathcal{G}(A) = \{f \oplus Af; f \in D(A)\}.$$

We set

$$\mathcal{G}'(A) = \{-Af \oplus f; f \in D(A)\}.$$

It is well-known that when A is densely defined closable operator, we have $\mathcal{G}(A)^\perp = \mathcal{G}'(A^*)$. We sometimes use the notation

$$\langle f, g \rangle_A = \langle f, g \rangle + \langle Af, Ag \rangle$$

$$(f, g)_A = (f, g) + (Af, Ag).$$

For $f, g \in L^2(0, \infty)$, we define $f \otimes g \in \mathbb{B}(L^2(0, \infty))$ by

$$(f \otimes g)h = (h, g)f, \quad h \in L^2(0, \infty).$$

We denote by $\mathbb{H}_r$ the right-half plane $\{z \in \mathbb{C}; \operatorname{Re} z > 0\}$. For $z \in \mathbb{H}_r$, we set $e_z(x) = e^{-zx}$. We denote by $L^1_{\text{loc}}[0, \infty)$ the set of measurable functions on $[0, \infty)$ that are integrable on every compact subset of $[0, \infty)$. For $f \in L^1_{\text{loc}}[0, \infty)$ with $a > 1$ such that $e_a f \in L^1(0, \infty)$, we denote by $\mathcal{L}[f](z)$ the Laplace transformation

$$\mathcal{L}[f](z) = \int_0^\infty f(x)e^{-zx}dx, \quad \operatorname{Re} z > a.$$

For $f \in L^2(0, \infty)$, we have $\mathcal{L}[f](z) = (f, e_z)$ and $|\mathcal{L}[f](z)| \leq \|f e_{z/2}\| \cdot \|e_{z/2}\|$, and so

$$(2.1) \quad \sqrt{x}|\mathcal{L}[f](x + iy)| \leq \|f e_{x/2}\| \rightarrow 0, \quad (x \rightarrow +\infty).$$

Let $S = \{S_t\}_{t \geq 0}$ be the shift semigroup of $L^2(0, \infty)$, that is

$$(S_t f)(x) = \begin{cases} 0 & (x < t) \\ f(x - t) & (t \leq x) \end{cases}.$$

The main purpose of this paper is to investigate the structure of C_0 -semigroups $T = \{T_t\}_{t \geq 0}$ of bounded operators in $\mathbb{B}(L^2(0, \infty))$ satisfying the following two conditions:

(C1) $T_t^* S_t = I$ for all $t \geq 0$.

(C2) $T_t - S_t$ is a Hilbert-Schmidt class operator for all $t \geq 0$.

Whenever a function $f \in L^2(0, \infty)$ extends to a continuous function on $[0, \infty)$, we abuse the notation and use the same symbol f for the extension. The generator A of $\{S_t\}_{t \geq 0}$ is the differential operator $(Af)(x) = -f'(x)$ with boundary condition $f(0) = 0$. More precisely, the domain $D(A)$ of A consists of square integrable locally absolutely continuous functions f on $[0, \infty)$ such that $f(0) = 0$ and the derivative f' (well-defined almost everywhere) belongs to $L^2(0, \infty)$. The adjoint operator A^* , which is the generator of $\{S_t^*\}_{t \geq 0}$, is the differential operator $(A^*f)(x) = f'(x)$ without any boundary condition. More precisely, $D(A^*)$ is the linear span of $D(A)$ and e^{-x} .

Let B be the generator of a semigroup $\{T_t\}_{t \geq 0}$ satisfying the condition (C1). Then for $f \in D(A)$ and $g \in D(B)$ we have $\langle S_t f, T_t g \rangle = \langle f, g \rangle$ and

$$\langle Af, g \rangle + \langle f, Bg \rangle = 0,$$

which shows $B \subset -A^*$. This means that B is also a differential operator, but what matters now is the domain of B .

For $p \in D(A^*) \setminus \{0\}$, we define $A_p \subset -A^*$ whose graph is $\mathcal{G}(-A^*) \cap \mathbb{C}(\bar{p} \oplus -\bar{p}')^\perp$, where $\bar{p}$ is the complex conjugate of p . By definition, A_p is a closed operator. As the codimension of $\mathcal{G}(A_p)$ in $\mathcal{G}(-A^*)$ is 1, the operator B is a restriction of A_p for some p (in fact, we will show $B = A_p$ later). Note that when $p(x) = e^{-x}$, the operator A_p is nothing but A .

Lemma 2.1. *The operator A_p is densely defined if and only if $p' \notin D(A) \setminus \{0\}$.*

Proof. Note that

$$D(A_p)^\perp \oplus 0 = (L^2(0, \infty) \oplus 0) \cap \mathcal{G}(A_p)^\perp = (L^2(0, \infty) \oplus 0) \cap \left(\mathcal{G}'(-A) + \mathbb{C}(\bar{p} \oplus -\bar{p}') \right).$$

Thus $g \in D(A_p)^\perp$ if and only if there exist $f \in D(A)$ and $\lambda \in \mathbb{C}$ such that

$$g = -f' + \lambda \bar{p},$$

$$0 = f - \lambda \bar{p}'.$$

This shows that if $p' \notin D(A)$, $g = 0$ and so the operator A_p is densely defined.

On the other hand, assume that $p' \in D(A) \setminus 0$. Then $\lambda(\bar{p} - p'')$ belongs to $D(A_p)^\perp$. Note that $p - p'' \neq 0$ as far as $p' \in D(A) \setminus \{0\}$. $\square$

We set

$$\mathcal{O} = \{p \in D(A^*); p' \notin D(A)\}.$$

Corollary 2.2. *Let $\{T_t\}_{t \geq 0}$ be a C_0 -semigroup satisfying (C1) and let B be the generator of $\{T_t\}_{t \geq 0}$. Then there exists $p \in \mathcal{O}$ such that $B \subset A_p$.*

We end this section with giving a proof for the claim stated in Introduction that the semigroup $\{T_t\}_{t \geq 0}$ satisfying (C1) and (C2) gives rise to an E_0 -semigroup. Let $H = e^{L^2(0, \infty)}$ be the symmetric Fock space for $L^2(0, \infty)$ and let Ω be the vacuum vector. We denote by $\exp(f) \in H$ the exponential vector for $f \in L^2(0, \infty)$, that is

$$\exp(f) = \bigoplus_{n=0}^{\infty} \frac{1}{\sqrt{n!}} f^{\otimes n},$$

where $f^{\otimes 0} = \Omega$. The Weyl operator $W(h)$ for $h \in L^2(0, \infty)$ is a unitary operator in $\mathbb{B}(H)$ determined by

$$W(h) \exp(f) = e^{-\frac{\|h\|^2}{2} - \langle f, h \rangle} \exp(f + h).$$

The CCR algebra for $L^2(0, \infty)$ is the algebra generated by $\{W(h)\}_{h \in L^2(0, \infty)}$ whose defining (vacuum) representation is irreducible.

Theorem 2.3. *If $\{T_t\}_{t \geq 0}$ satisfies (C1) and (C2) and T_t preserves the real functions $L^2(0, \infty)_{\mathbb{R}}$, then there exists a unique E_0 -semigroup $\{\alpha_t\}_{t \geq 0}$ acting on $\mathbb{B}(H)$ satisfying.*

$$\alpha_t(W(f + ig)) = W(S_t f + i T_t g), \quad f, g \in L^2(0, \infty)_{\mathbb{R}}.$$

Proof. The condition (C1) shows that there exists a representation β_t of the CCR algebra for $L^2(0, \infty)$ satisfying $\beta_t(W(f + ig)) = W(S_t f + iT_t g)$ for $f, g \in L^2(0, \infty)_{\mathbb{R}}$. To show the claim, it suffices to prove that β_t is quasi-equivalent to the vacuum representation. We set M to be the von Neumann algebra generated by the image of β_t . Since M is a factor thanks to [1, Theorem 1], it suffices to show that the representation β_t restricted to $\overline{M\Omega}$ is quasi-equivalent to the vacuum representation. Thus our problem is now reduced to the quasi-equivalence of the vacuum representation and the GNS representation of the quasi-free state $\omega'(x) = \langle \beta_t(x)\Omega, \Omega \rangle$, which can be shown by using (C2) from a well-known criterion (see [2],[5] for example). $\square$

3. RESOLVENT

In this section, we explicitly compute the resolvent of the generator of a semigroup $\{T_t\}_{t \geq 0}$ satisfying (C1). It turns out that such a semigroup is completely characterized by a holomorphic function on the right-half plane.

For $p \in D(A^*)$, we set

$$\begin{aligned} \mathcal{M}[p](z) &:= (p, e_z)_{A^*} = \mathcal{L}[p](z) - z\mathcal{L}[p'](z) = (1 - z^2)\mathcal{L}[p](z) + p(0)z \\ &= \left(\frac{1}{z} - z\right)\mathcal{L}[p'](z) + \frac{p(0)}{z}. \end{aligned}$$

Note that unless $p = 0$, the function $\mathcal{M}[p](z)$ is not constantly zero.

For a point $z \in \mathbb{H}_r$ with $\mathcal{M}[p](z) \neq 0$, we define $\xi_{p,z}(x)$ by

$$\begin{aligned} \xi_{p,z}(x) &= -\frac{\int_0^\infty e^{-zt} \left(p(x+t) + z(p'(x) - p'(x+t)) \right) dt}{\mathcal{M}[p](z)} \\ &= -\frac{(1 - z^2) \int_0^\infty e^{-zt} p(x+t) dt + zp(x) + p'(x)}{\mathcal{M}[p](z)}. \end{aligned}$$

Note that $\xi_{p,z}$ is given by the $L^2(0, \infty)$ -valued integral

$$\xi_{p,z} = -\frac{(1 - z^2) \int_0^\infty e^{-zt} S_t^* p dt + zp + p'}{\mathcal{M}[p](z)}$$

and so $\xi_{p,z} \in L^2(0, \infty)$.

Lemma 3.1. *Let $p \in \mathcal{O}$ and $\operatorname{Re} z > 0$. Then z is in the resolvent set of A_p if and only if $\mathcal{M}[p](z) \neq 0$. If z satisfies this condition, the resolvent $(zI - A_p)^{-1}$ is given by*

$$(zI - A_p)^{-1} = (zI - A)^{-1} + e_z \otimes \xi_{p,z}.$$

Proof. If $\mathcal{M}[p](z) = 0$, we have $e_z \in D(A_p)$ and $A_p e_z = z e_z$. Thus $\mathcal{M}[p](z) \neq 0$ is a necessary condition for z to belong to the resolvent set of A_p .

Assume $\mathcal{M}[p](z) \neq 0$ now. Let

$$B_z = (zI - A)^{-1} + e_z \otimes \xi_{p,z}.$$

Note that z belongs to the resolvent set of A for $\operatorname{Re} z > 0$ and the resolvent is given by

$$(zI - A)^{-1}f(x) = \int_0^x f(t)e^{-z(x-t)}dt.$$

First we show $B_z(zI - A_p)g = g$ for $g \in D(A_p)$. As

$$(zI - A)^{-1}(zI - A_p)g(x) = \int_0^x (zg(t) + g'(t))e^{-z(x-t)}dt = g(x) - g(0)e^{-zx},$$

it suffices to show $(zI - A_p)g, \xi_{p,z}) = g(0)$. Indeed,

$$\begin{aligned} ((zI - A_p)g, \xi_{p,z}) &= -\frac{\int_0^\infty e^{-zt}((zI - A_p)g, S_t^*(p - zp'))dt + (zg + g', p')}{\mathcal{M}[p](z)} \\ &= -\frac{\int_0^\infty e^{-zt}(S_t(zI - A_p)g, p - zp')dt + (zg + g', p')}{\mathcal{M}[p](z)} \\ &= -\frac{((zI - A)^{-1}(zI - A_p)g, p - zp') + (zg + g', p')}{\mathcal{M}[p](z)} \\ &= -\frac{(g - g(0)e_z, p - zp') + (zg + g', p')}{\mathcal{M}[p](z)} \\ &= -\frac{(g, p) + (g', p') - g(0)((e_z, p) - z(e_z, p'))}{\mathcal{M}[p](z)} \\ &= g(0). \end{aligned}$$

Next we show that the range of B_z is contained in $D(A_p)$. Note that for $f \in L^2(0, \infty)$ the element $B_z f = (zI - A)^{-1}f + (f, \xi_{p,z})e_z$ belongs to $D(A^*)$, and so

$$\begin{aligned} (B_z f, p) + (A^* B_z f, A^* p) &= ((zI - A)^{-1}f, p) + (A^*(zI - A)^{-1}f, A^* p) + ((e_z, p) + (A^* e_z, A^* p))(f, \xi_{p,z}) \\ &= ((zI - A)^{-1}f, p) - (A(zI - A)^{-1}f, A^* p) + \mathcal{M}[p](z)(f, \xi_{p,z}) \\ &= ((zI - A)^{-1}f, p) + (f - z(zI - A)^{-1}f, A^* p) + \mathcal{M}[p](z)(f, \xi_{p,z}) \\ &= ((zI - A)^{-1}f, p) + (f, A^* p) - z(A(zI - A)^{-1}f, p) + \mathcal{M}[p](z)(f, \xi_{p,z}) \\ &= (1 - z^2)((zI - A)^{-1}f, p) + (f, p') + z(f, p) + \mathcal{M}[p](z)(f, \xi_{p,z}) \\ &= (1 - z^2)((zI - A)^{-1}f, p) - (1 - z^2) \int_0^\infty e^{-zt}(f, S_t^* p) \\ &= (1 - z^2)((zI - A)^{-1}f, p) - (1 - z^2) \int_0^\infty e^{-zt}(S_t f, p) \\ &= 0. \end{aligned}$$

Thus, we get $B_z f \in D(A_p)$ and

$$\begin{aligned} (zI - A_p)B_z f &= (zI + A^*)((zI - A)^{-1}f + (f, \xi_{p,z})e_z) \\ &= (zI + A^*)(zI - A)^{-1}f = f. \end{aligned}$$

Therefore, $B_z = (zI - A_p)^{-1}$. □

Note that when A_1 and A_2 are densely defined closed operators on a Banach space such that A_1 is a proper extension of A_2 , the intersection of the resolvent sets of A_1 and A_2 is empty.

Theorem 3.2. *Let $\{T_t\}_{t \geq 0}$ be a C_0 -semigroup satisfying (C1). Then there exists $p \in \mathcal{O}$ such that A_p is the generator of $\{T_t\}_{t \geq 0}$. Moreover, there exists a positive number a such that for $\forall f \in D(A_p)$, $\forall t > 0$ and $\forall b > a$, the function $T_t f$ is given by*

$$T_t f = S_t f + \lim_{r \rightarrow +\infty} \frac{1}{2\pi i} \int_{b-ir}^{b+ir} e^{zt} (f, \xi_{p,z}) e_z dz.$$

Proof. Let B be a generator of $\{T_t\}_{t \geq 0}$. Thanks to Corollary 2.2, there exists $p \in D(A^*)$ with $p' \notin D(A)$ such that $B \subset A_p$. It is well-known that there exist positive numbers a and M such that for all $t \geq 0$ the inequality $\|T_t\| \leq M e^{at}$ holds [19, p.232]. Thus z belongs to the resolvent set of B for $\operatorname{Re} z > a$ and

$$(zI - B)^{-1} f = \int_0^\infty e^{-zt} T_t f, \quad \forall f \in L^2(0, \infty).$$

On the other hand, since $\mathcal{M}[p](z)$ is a non-constant analytic function on $\mathbb{H}_r$, there exists z_0 such that $\operatorname{Re} z_0 > a$ and $\mathcal{M}[p](z_0) \neq 0$, and so Lemma 3.2 implies that z_0 is in the resolvent set of A_p . If B were proper restriction of A_p , the point z_0 would not belong to the resolvent set of B , which is a contradiction. Thus $B = A_p$. The integral formula above follows from the usual inverse Laplace transformation with the fact that the map $(0, \infty) \ni t \mapsto T_t f \in L^2(0, \infty)$ is differentiable for $f \in D(A_p)$. □

The above theorem shows that every information of the semigroup $\{T_t\}_{t \geq 0}$ is encoded in $\xi_{p,z}$, which is determined by $\mathcal{M}[p]$ via the following lemma:

Lemma 3.3. *Let $p \in \mathcal{O}$ and let z satisfy $\operatorname{Re} z > 0$ and $\mathcal{M}[p](z) \neq 0$. Then*

$$\mathcal{L}[\xi_{p,z}](w) = \frac{\mathcal{M}[p](z) - \mathcal{M}[p](w)}{\mathcal{M}[p](z)(z - w)}.$$

Proof. Assume $w \neq z$. Then

$$\begin{aligned}
& \mathcal{M}[p](z)\mathcal{L}[\xi_{p,z}](w) \\
&= -\mathcal{L}[p'](w) - z\mathcal{L}[p](w) - (1-z^2) \int_0^\infty dt \int_0^\infty dx e^{-wx} e^{-zt} p(x+t) \\
&= p(0) - (z+w)\mathcal{L}[p](w) - (1-z^2) \int_0^\infty dt \int_t^\infty dx e^{-w(x-t)} e^{-zt} p(x) \\
&= p(0) - (z+w)\mathcal{L}[p](w) - (1-z^2) \int_0^\infty p(x) e^{-wx} dx \int_0^x dt e^{-(z-w)t} \\
&= p(0) - (z+w)\mathcal{L}[p](w) + (1-z^2) \int_0^\infty p(x) \frac{e^{-zx} - e^{-wx}}{z-w} dx \\
&= p(0) - (z+w)\mathcal{L}[p](w) + (1-z^2) \frac{\mathcal{L}[p](z) - \mathcal{L}[p](w)}{z-w} \\
&= \frac{\mathcal{M}[p](z) - \mathcal{M}[p](w)}{z-w},
\end{aligned}$$

which shows the statement. $\square$

When A_p generates a C_0 -semigroup $\{T_t\}_{t>0}$, we call p the *orthogonal function* of $\{T_t\}_{t>0}$ and call $\mathcal{M}[p]$ the *half density function* of $\{T_t\}_{t>0}$. The half density function plays the central role throughout this note and the theory of the Hardy classes is best suit for its description.

We denote by $H^p(\mathbb{H}_r)$ the Hardy space of the right half plane $\mathbb{H}_r$, that is, the set of holomorphic functions F on $\mathbb{H}_r$ such that $\sup_{x>0} \|F(x+i\cdot)\|_p$ is finite. For $F \in H^p(\mathbb{H}_r)$ and iy in the imaginary axis, we denote by $F(iy)$ the non-tangential limit of F at iy , which makes sense for almost every iy . Note that when $f \in L^2(0, \infty)$, its Laplace transform $\mathcal{L}[f]$ belongs to $H^2(\mathbb{H}_r)$. Thanks to the Paley-Wiener theorem, every element of $H^2(\mathbb{H}_r)$ is of this form.

Let $\mathcal{HD}$ be the set of holomorphic functions $M(z)$ on $\mathbb{H}_r$ such that $M(z)/(1+z)$ belongs to $H^2(\mathbb{H}_r)$ and M does not belong to $H^2(\mathbb{H}_r)$.

Lemma 3.4. *Let the notation be as above. Then,*

- (1) *The map $p \mapsto \mathcal{M}[p]$ gives a bijection between $\mathcal{O}$ and $\mathcal{HD}$.*
- (2) *The map $q \mapsto (1+z)\mathcal{L}[q](z)$ gives a bijection between $L^2(0, \infty) \setminus D(A)$ and $\mathcal{HD}$.*
- (3) *The bijection between $\mathcal{O}$ and $L^2(0, \infty) \setminus D(A)$ induced by (1) and (2) is given by*

$$\begin{aligned}
q(x) &= p(x) - p'(x) - p(0)e^{-x}, \\
p(x) &= 2\mathcal{L}[q](1) \cosh x - \int_0^x e^{x-t} q(t) dt = \mathcal{L}[q](1) e^{-x} + \int_0^\infty e^{-t} S_t^* q(x) dt.
\end{aligned}$$

Proof. (2) Since $D(A)$ is exactly the set of $q \in L^2(0, \infty)$ with $z\mathcal{L}[q](z) \in H^2(\mathbb{H}_r)$, this is a direct consequence of the the Paley-Wiener theorem.

(1) Let $p \in \mathcal{O}$. Using $\mathcal{L}[p'](x) = z\mathcal{L}[p](z) - p(0)$, we get

$$\frac{\mathcal{M}[p](z)}{1+z} = \mathcal{L}[p](z) - \mathcal{L}[p'](z) - \frac{p(0)}{1+z} = \mathcal{L}[p - p' - p(0)e_1](z) \in H^2(\mathbb{H}_r).$$

Assume that $\mathcal{M}[p](z)$ belongs to $H^2(\mathbb{H}_r)$. Then since we have

$$\frac{z\mathcal{M}[p](z)}{1+z} = (1-z)\mathcal{L}[p'](z) + \frac{p(0)}{1+z},$$

we would get

$$z\mathcal{L}[p'](z) = \mathcal{L}[p'](z) - \frac{z\mathcal{M}[p](z)}{1+z} + \frac{p(0)}{1+z} \in H^2(\mathbb{H}_r).$$

This implies $p' \in D(A)$, which is contradiction. Thus $\mathcal{M}[p] \in \mathcal{HD}$.

Injectivity of the map $p \mapsto \mathcal{M}[p]$ is obvious and we show surjectivity now. For $M \in \mathcal{HD}$, take $q \in L^2(0, \infty) \setminus D(A)$ satisfying $M(z) = (1+z)\mathcal{L}[q](z)$. We define $p(x)$ by the second formula in (3). Then p is an absolutely continuous function in $L^2(0, \infty)$ satisfying the first formula and so $p \in D(A^*)$. Since $p' - q = p - p(0)e_1 \in D(A)$ and q does not belong to $D(A)$, we conclude $p' \notin D(A)$ and so $p \in \mathcal{O}$. $M = \mathcal{M}[p]$ can be shown by computing the Laplace transform.

(3) has already been shown in the above argument. $\square$

From now on, we are mainly working on $\mathcal{HD}$ and $L^2(0, \infty) \setminus D(A)$ instead of $\mathcal{O}$. When $p \in \mathcal{O}$ and $M = \mathcal{M}[p]$, we abuse the notation and use the symbols A_M and $\xi_{M,z}$ for A_p and $\xi_{p,z}$ respectively. Lemma 3.3 means

$$(3.1) \quad \mathcal{L}[\xi_{M,z}](w) = \frac{M(z) - M(w)}{M(z)(z-w)}.$$

When $(1+z)\mathcal{L}[q](z) = M(z)$, we can show the following by computing the Laplace transform:

$$(3.2) \quad \xi_{M,z}(y) = \frac{q(y) - (1+z) \int_0^\infty q(y+s)e^{-sz}ds}{M(z)}.$$

We denote by $\mathcal{HD}_b$ the set of $M \in \mathcal{HD}$ such that A_M generates a C_0 -semigroup $\{T_t\}_{t>0}$. Such $\{T_t\}_{t>0}$ always satisfies (C1). We denote by $\mathcal{HD}_2$ the set of $M \in \mathcal{HD}_b$ such that $\{T_t\}_{t>0}$ satisfies (C1) and (C2).

When $x+iy \in \mathbb{H}_r$ and f is a measurable function on $\mathbb{R}$ such that $f(\lambda)/(1+\lambda^2) \in L^1(\mathbb{R})$, we denote by $\mathcal{P}[f](x+iy)$ the Poisson integral

$$\mathcal{P}[f](x+iy) = \frac{1}{\pi} \int_{-\infty}^\infty \frac{xf(\lambda)}{x^2 + (y-\lambda)^2} d\lambda.$$

Lemma 3.5. *Let $M \in \mathcal{HD}$ and assume $z = x+iy \in \mathbb{H}_r$ is not a zero of M . Then*

$$|\xi_{M,z}|^2 = \frac{1}{2\operatorname{Re} z} \left(\frac{\mathcal{P}[|M(i\cdot)|^2](z)}{|M(z)|^2} - 1 \right).$$

Proof. The Plancherel theorem implies

$$\begin{aligned}
\|\xi_{M,z}\|^2 &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} |\mathcal{L}[\xi_{M,z}](i\lambda)|^2 d\lambda \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{|M(z) - M(i\lambda)|^2}{|M(z)(z - i\lambda)|^2} d\lambda \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{|M(z)|^2 - 2\operatorname{Re}(M(i\lambda)\overline{M(z)}) + |M(i\lambda)|^2}{|M(z)|^2|z - i\lambda|^2} d\lambda \\
&= \frac{1}{2\operatorname{Re} z |M(z)|^2} [|M(z)|^2 - 2\operatorname{Re}(\mathcal{P}[M(i\cdot)](z)\overline{M(z)}) + \mathcal{P}[|M(i\cdot)|^2](z)].
\end{aligned}$$

Thus it suffices to show $\mathcal{P}[M(i\cdot)](z) = M(z)$. Let $re^{i\theta} = (z-1)/(z+1)$ and set $f(re^{i\theta}) = M(z)$. Since $M(z)/(1+z) \in H^2(\mathbb{H}_r)$, the function f belongs to the Hardy space $H^2(\mathbb{D})$ of the unit disc $\mathbb{D}$ [8, Chapter 8] and

$$\mathcal{P}[M(i\cdot)](z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{(1-r^2)f(e^{it})}{1-2r\cos(\theta-t)+r^2} dt.$$

Therefore $\mathcal{P}[M(i\cdot)](z) = M(z)$ follows from the usual Poisson integral formula. $\square$

Note that $\mathcal{P}[|M(i\cdot)|^2](z) \geq |M(z)|^2$ always holds.

Corollary 3.6. *Let $M \in \mathcal{HD}$. If A_M generates a C_0 -semigroup, then there exist positive constants a and C such that for all $\operatorname{Re} z \geq a$,*

$$\log \mathcal{P}[|M(i\cdot)|^2](z) - \log |M(z)|^2 \leq \frac{\mathcal{P}[|M(i\cdot)|^2](z)}{|M(z)|^2} - 1 \leq C.$$

Moreover,

$$\sqrt{x}|M(x)| \rightarrow +\infty \quad (x \rightarrow +\infty).$$

Proof. Lemma 3.1 shows that if $M \in \mathcal{HD}$ and $M(z) \neq 0$,

$$\begin{aligned}
\frac{\|\xi_{M,z}\|}{\sqrt{2\operatorname{Re} z}} &= \|e_z \otimes \xi_{M,z}\| = \|(zI - A_h)^{-1} - (zI - A)^{-1}\| \\
&\leq \|(zI - A_h)^{-1}\| + \|(zI - A)^{-1}\|.
\end{aligned}$$

When A_M generates a C_0 -semigroup, there exist constant $a > 0$ and C_1 such that $\|(zI - A_M)^{-1}\| \leq C_1/\operatorname{Re} z$ for all $\operatorname{Re} z > a$. Thus the first statement follows from Lemma 3.5. Moreover, $\{x(xI - A_M)^{-1}\}_{x \geq a}$ converges to 0 in the strong topology as x tends to $+\infty$. Thanks to Lemma 3.1 and Lemma 3.3, for $w \in \mathbb{H}_r$ we have

$$\begin{aligned}
\|x(xI - A_M)^{-1}e_w - x(xI - A)^{-1}e_w\| &= x|(\xi_{M,x}, e_w)| \|e_x\| \\
&= \sqrt{\frac{1}{2x}} \frac{|1 - \frac{M(w)}{M(x)}|}{|1 - \frac{w}{x}|} \rightarrow 0, \quad (x \rightarrow +\infty),
\end{aligned}$$

which shows the statement. $\square$

4. GLOBAL THEORY

Let $\{S_t\}_{t \geq 0}$ be as before and $\{K_t\}_{t \geq 0}$ be a family of bounded operators on $L^2(0, \infty)$. Then it is easy to show that the C_0 -semigroup condition for $\{T_t = S_t + K_t\}$ is equivalent to (K0) and that the condition (C1) is equivalent to (K1) below respectively:

(K0) The map $\mathbb{R}_{>0} \ni t \mapsto K_t$ is strongly continuous and

$$K_{s+t} = K_s S_t + S_s K_t + K_s K_t, \quad \forall s, t \geq 0,$$

(K1)

$$K_t L^2(0, \infty) \subset L^2(0, t), \quad \forall t \geq 0.$$

We claim that the above conditions and (C2) altogether are equivalent to

(K2) There exists a measurable function $k(x, y)$ defined on $(0, \infty)^2$ satisfying

$$K_t f(x) = \begin{cases} \int_0^t k(t-x, y) f(y) dy & (x < t) \\ 0 & (t \leq x) \end{cases},$$

$$k(x+t, y) = k(x, y+t) + \int_0^t k(x, s) k(t-s, y) ds, \quad \forall t \geq 0, \text{ a.e. } (x, y) \in (0, \infty)^2,$$

$$\int_0^t \int_0^\infty |k(x, y)|^2 dy dx < +\infty, \quad \forall t > 0.$$

Indeed, assume that $\{T_t\}_{t \geq 0}$ is a C_0 -semigroup satisfying (C1) and (C2). Let k_t be the integral kernel for K_t . Then $k_t \in L^2(0, \infty)^2$ with support in $(0, t] \times (0, \infty)$. The semigroup property of $\{T_t\}$ implies that for every $s, t \geq 0$ and $f \in L^2(0, \infty)$, the following holds:

$$\begin{aligned} & \int_0^\infty k_{s+t}(x, y) f(x) dx \\ &= \int_t^\infty k_s(x, y) f(y-t) dy + 1_{[s, s+t]}(x) \int_0^\infty k_t(x-s, y) f(y) dy \\ &+ \int_0^\infty \int_0^t k_s(x, u) k_t(u, y) f(y) du dy \\ &= \int_0^\infty \left(k_s(x, y+t) + 1_{[s, s+t]}(x) k_t(x-s, y) + \int_0^t k_s(x, u) k_t(u, y) du \right) f(y) dy, \end{aligned}$$

where $1_I(x)$ denotes the characteristic function of an interval I . Therefore, for almost all $(x, y) \in (0, \infty)^2$, we get

$$k_{s+t}(x, y) = k_s(x, y+t) + 1_{[s, s+t]}(x) k_t(x-s, y) + \int_0^t k_s(x, u) k_t(u, y) du.$$

We set $\tilde{k}_t(x, y) = k_t(t-x, y)$ for $x \in (0, t]$ and set $\tilde{k}_t(x, y) = 0$ for $x > t$. Then the above relation is equivalent to

$$\begin{aligned} \tilde{k}_{s+t}(s+t-x, y) &= 1_{(0, s]}(x) \left(\tilde{k}_s(s-x, y+t) + \int_0^t \tilde{k}_s(s-x, u) \tilde{k}_t(t-u, y) du \right) \\ &+ 1_{[s, s+t]}(x) \tilde{k}_t(s+t-x, y), \end{aligned}$$

for $0 \leq x \leq s + t$. This shows that there exists a measurable function $k(x, y)$ on $(0, 1)^2$ such that $\tilde{k}_t(x, y) = k(x, y)$ for $(x, y) \in (0, t] \times (0, \infty)$ and (K2) holds. The converse implication also follows from the same computation.

Lemma 4.1. *Let $\{T_t\}_{t \geq 0}$ be a C_0 -semigroup satisfying (C1) and (C2) and let K_t and $k(x, y)$ be as above. We define an $L^2(0, \infty)$ -valued function $\kappa(x)$ on $(0, \infty)$ by $\kappa(x)(\cdot) = k(x, \cdot)$. Then*

- (1) *There exists a positive number a such that $e^{-ax}k(x, y)$ is square integrable on $(0, \infty)^2$. In particular,*

$$\mathcal{L}[\kappa](z)(\cdot) := \int_0^\infty e^{-zx}k(x, \cdot)dx$$

makes sense as $L^2(0, \infty)$ -valued holomorphic function on $\operatorname{Re} z > a$.

- (2) *Let A_M be the generator of $\{T_t\}_{t \geq 0}$ with $M \in \mathcal{HD}$. Then $\xi_{M,z} = \mathcal{L}[\kappa](z)$ and*

$$\int_0^\infty \int_0^\infty |e^{-rx}k(x, y)|^2 dx dy = \frac{1}{2\pi} \int_{-\infty}^\infty \|\xi_{M,r+is}\|^2 ds, \quad \forall r \geq a.$$

Proof. (1) The condition (K0) implies

$$\|K_{s+t}\|_{H.S.} \leq \|K_s\|_{H.S.} + \|K_t\|_{H.S.} + \|K_s\|_{H.S.}\|K_t\|,$$

where $\|\cdot\|_{H.S.}$ denotes the Hilbert-Schmidt norm. Thus for every natural number $n \in \mathbb{N}$, we get

$$\|K_{n+1}\|_{H.S.} \leq \|K_n\|_{H.S.} + \|K_1\|_{H.S.} + \|K_n\|_{H.S.}\|K_1\|,$$

which implies

$$\|K_n\|_{H.S.} \leq \frac{\|K_1\|_{H.S.}}{\|K_1\|} (1 + \|K_1\|)^{n+1}.$$

Since $\|K_t\|_{H.S.} = \left(\int_0^t \int_0^\infty |k(x, y)|^2 dy dx \right)^{1/2}$ is an increasing function in t with the above estimate for $t = n \in \mathbb{N}$, there exist two positive numbers M and b such that

$$\|K_t\|_{H.S.}^2 \leq M e^{bt}.$$

We choose a satisfying $2a > b$ and set $\theta(t) = \|K_t\|_{H.S.}^2$. Then

$$\begin{aligned} \int_0^r \int_0^\infty |e^{-ax}k(x, y)|^2 dy dx &= \int_0^r e^{-2at} \theta'(t) dt = \theta(r) e^{-2ar} + 2a \int_0^r \theta(t) e^{-2at} dt \\ &\leq M e^{-(2a-b)r} + 2a \int_0^r M e^{-(2a-b)t} dt \\ &\leq \frac{2aM}{2a-b}, \end{aligned}$$

which shows $e^{-ax}k(x, y)$ is square integrable.

(2) We compute the Laplace transformation of K_t . For $f, g \in L^2(0, \infty)$ and z with $\operatorname{Re} z > a$, we get

$$\begin{aligned} \int_0^\infty e^{-zt} (K_t f, g) dt &= \int_0^\infty \int_0^t \int_0^\infty e^{-zt} k(t-x, y) f(y) g(x) dy dx dt \\ &= \int_0^\infty \int_x^\infty \int_0^\infty e^{-zt} k(t-x, y) f(y) g(x) dy dt dx \\ &= \int_0^\infty \int_0^\infty \int_0^\infty e^{-z(t+x)} k(t, y) f(y) g(x) dy dt dx \\ &= (\mathcal{L}[\kappa](z), f)(e_z, g). \end{aligned}$$

Therefore Lemma 3.1 and Theorem 3.2 imply $\xi_{M,z} = \mathcal{L}[\kappa](z)$. The last equation follows from the Plancherel theorem. $\square$

Theorem 4.2. *Let $M \in \mathcal{HD}$. Then $M \in \mathcal{HD}_2$ if and only if there exist positive numbers a and C such that $\xi_{M,z}$ is holomorphic on $\operatorname{Re} z > a$ and for all $x > a$*

$$\int_{-\infty}^{+\infty} \|\xi_{M,x+iy}\|^2 dy \leq C.$$

Proof. Assume that A_M generates a C_0 -semigroup satisfying the conditions (C1) and (C2). Lemma 3.1 and Lemma 4.1 imply that there exist positive numbers a and M satisfying the above property.

Assume now that there exist a and M satisfying the above property. Then Paley-Wiener theorem [19, p. 163] implies that there exists a $L^2(0, \infty)$ -valued measurable function $\kappa(x)$ on $(0, \infty)$ such that

$$\xi_{M,z} = \int_0^\infty \kappa(x) e^{-zx} dx$$

and $e^{-ax} \|\kappa(x)\|$ is square integrable.

We define a family of Hilbert-Schmidt operators $K_t \in \mathbb{B}(L^2(0, \infty))$ for $t \geq 0$ by

$$K_t f(x) = \begin{cases} (\kappa(t-x), f) & (x < t) \\ 0 & (t \leq x) \end{cases}$$

Note that

$$\int_0^\infty e^{-zt} K_t f = (f, \xi_{M,z}) e_z$$

holds for all $f \in L^2(0, \infty)$ and z with $\operatorname{Re} z > a$. It is easy to show that the map $t \mapsto K_t$ is continuous in the strong operator topology. Thus to show that $\{S_t + K_t\}_{t \geq 0}$ is a C_0 -semigroup generated by A_M , it suffices to prove

$$K_{s+t} = K_s S_t + S_s K_t + K_s K_t, \quad \forall s, t \geq 0,$$

or equivalently to prove

$$\begin{aligned} \int_0^\infty ds \int_0^\infty dt e^{-(z_1 s + z_2 t)} (K_{s+t} f, g) \\ = \int_0^\infty ds \int_0^\infty dt e^{-(z_1 s + z_2 t)} (K_s S_t f + S_s K_t f + K_s K_t f, g) \end{aligned}$$

for all f, g in a total set in $L^2(0, \infty)$ and for all $\operatorname{Re} z_1 > a$, $\operatorname{Re} z_2 > a$. Assume $z_1 \neq z_2$. Then the left-hand side is

$$\begin{aligned} & \int_0^\infty ds \int_0^\infty dt e^{-(z_1 s + z_2 t)} (K_{s+t} f, g) = \int_0^\infty ds \int_s^\infty dt e^{-(z_1 s + z_2(t-s))} (K_t f, g) \\ &= \int_0^\infty dt (K_t f, g) e^{-z_2 t} \int_0^t ds e^{-(z_1 - z_2)s} = \int_0^\infty dt (K_t f, g) \frac{e^{-z_2 t} - e^{-z_1 t}}{z_1 - z_2} \\ &= \frac{(f, \xi_{M, z_2})(e_{z_2}, g) - (f, \xi_{M, z_1})(e_{z_1}, g)}{z_1 - z_2}. \end{aligned}$$

The right-hand side is

$$\begin{aligned} & ((z_2 I - A)^{-1} f, \xi_{M, z_1})(e_{z_1}, g) + (f, \xi_{M, z_2})((z_1 I - A)^{-1} e_{z_2}, g) \\ &+ (f, \xi_{M, z_2})(e_{z_2}, \xi_{M, z_1})(e_{z_1}, g) \\ &= ((z_2 I - A)^{-1} f, \xi_{M, z_1})(e_{z_1}, g) + \frac{(f, \xi_{M, z_2})(e_{z_2}, g) - (f, \xi_{M, z_2})(e_{z_1}, g)}{z_1 - z_2} \\ &+ (f, \xi_{M, z_2})(e_{z_1}, g) \frac{M(z_1) - M(z_2)}{(z_1 - z_2)M(z_1)} \\ &= ((z_2 I - A)^{-1} f, \xi_{M, z_1})(e_{z_1}, g) + \frac{(f, \xi_{M, z_2})(e_{z_2}, g)}{z_1 - z_2} - \frac{(f, \xi_{M, z_2})(e_{z_1}, g)M(z_2)}{(z_1 - z_2)M(z_1)}, \end{aligned}$$

where we use $(\xi_{M, z}, e_w) = \mathcal{L}[\xi_{M, z}](w)$ and Lemma 3.3. Thus, it suffices to show

$$((z_2 I - A)^{-1} f, \xi_{M, z_1}) = \frac{(f, \xi_{M, z_2})M(z_2) - (f, \xi_{M, z_1})M(z_1)}{(z_1 - z_2)M(z_1)}.$$

Setting $f = e_w$ with $\operatorname{Re} w > 0$ and $w \neq z_1$, $w \neq z_2$, we can show this equality using Lemma 3.3, and so we get the statement. $\square$

Corollary 4.3. *Let $M(z)$ be a holomorphic function on $\mathbb{H}_r$.*

(1) *Assume that $M \in \mathcal{HD}_2$. then*

(i)

$$\sup_{y \in \mathbb{R}} \frac{\mathcal{P}[|M(i \cdot)|^2](x + iy)}{|M(x + iy)|^2} = O(1), \quad (x \rightarrow +\infty),$$

(ii)

$$\int_{-\infty}^{+\infty} \left(\frac{\mathcal{P}[|M(i \cdot)|^2](x + iy)}{|M(x + iy)|^2} - 1 \right) dy = o(x), \quad (x \rightarrow +\infty).$$

(iii) *There exists positive constant $a > 0$ and $f \in L^2_{\text{loc}}(0, \infty)$ such that $e_a f \in L^2(0, \infty)$ and $1/(zM(z))$ is the Laplace transformation of f . In particular, there exists a positive constant C such that*

$$|M(z)| \geq \frac{C \sqrt{\operatorname{Re} z - a}}{|z|}, \quad \operatorname{Re} z > a.$$

(2) If M satisfies $M(z)/(1+z) \in H^2(\mathbb{H}_r)$ and

$$\int_{-\infty}^{+\infty} \left(\frac{\mathcal{P}[|M(i\cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right) dy = O(x), \quad (x \rightarrow +\infty),$$

then $M \in \mathcal{HD}_2$.

Proof. (1) (i) follows from Corollary 3.6.

Let the notation be as in the proof of the previous theorem. Then

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \|\xi_{M,x+iy}\|^2 dy = \int_0^{\infty} \|\kappa(s)\|^2 e^{-2xs} ds \rightarrow 0, \quad (x \rightarrow +\infty),$$

which together with Lemma 3.5 and Lemma 4.1 implies (ii).

Take $a > 0$ such that

$$\int_0^{\infty} \int_0^{\infty} |e^{-as} k(s, t)|^2 ds dt < \infty.$$

For $x \geq a$, we define a Hilbert-Schmidt operator L_x by

$$L_x f(s) = \int_0^{\infty} e^{-xs} k(s, t) f(t) dy.$$

Lemma 3.3 and Lemma 4.1 imply

$$\mathcal{L}[L_x e_w](z) = \frac{M(x+z) - M(w)}{(x+z-w)M(x+z)},$$

and so,

$$\|L_x e_w\|^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{1 - \frac{M(w)}{M(x+iy)}}{x+iy-w} \right|^2 dy.$$

On the other hand,

$$\|L_x e_w\| \leq \|L_x\| \|e_w\| \leq \|L_a\|_{H.S.} \|e_w\|,$$

and we get

$$\sup_{x \geq a} \int_{-\infty}^{\infty} \frac{dy}{|x+iy|^2 |M(x+iy)|^2} < \infty.$$

Thus (iii) follows from the Paley-Winer theorem.

(2) Let M be a function satisfying the assumption. If $M \notin H^2(\mathbb{H}_r)$, we are done thanks to Theorem 4.2. Note that $\xi_{M,z}$ still makes sense for M without the condition $M \notin H^2(\mathbb{H}_r)$. Using the same argument as in the proof of Theorem 4.2, we can construct a C_0 -semigroup $\{T_t\}_{t>0}$, whose Laplace transformation is given by $(zI - A)^{-1} + e_z \otimes \xi_{M,z}$. Let A_{M_1} be the generator of the $\{T_t\}_{t>0}$. Then we have $\xi_{M,z} = \xi_{M_1,z}$, and so

$$\frac{M(z) - M(w)}{M(z)(z-w)} = \frac{M_1(z) - M_1(w)}{M_1(z)(z-w)}$$

thanks to Lemma 3.3. Letting w tend to z , we get $M'(z)/M(z) = M'_1(z)/M_1(z)$, and consequently M is a scalar multiple of M_1 , which shows $M \notin H^2(\mathbb{H}_r)$. $\square$

Corollary 4.4. *Let $M(z)$ be a holomorphic function on $\mathbb{H}_r$ satisfying $M(z)/(1+z) \in H^2(\mathbb{H}_r)$. We assume that there exist positive constants a, n , and C such that*

$$\frac{1}{|M(z)|} \leq C(1+|z|)^n$$

holds for all z with $\operatorname{Re} z \geq a$ and

$$\int_{-\infty}^{+\infty} \left(\frac{\mathcal{P}[|M(i\cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right) dy < +\infty$$

for some $a < x$. Then $M \in \mathcal{HD}_2$.

Proof. First we claim that there exists a distribution $k \in \mathcal{D}(\mathbb{R}^2)'$ whose support is in $[0, \infty)^2$ such that the function $\mathcal{L}[\xi_{M,z}](w)$ in (z, w) is the Laplace transformation of $k(s, t)$. Indeed, let $q \in L^2(0, \infty)$ satisfying $(1+z)\mathcal{L}[q](z) = M(z)$. Lemma 3.3 shows

$$M(z)\mathcal{L}[\xi_{M,z}](w) = (1+z) \frac{\mathcal{L}[q](z) - \mathcal{L}[q](w)}{z-w} + \mathcal{L}[q](w).$$

Note that

$$-\frac{\mathcal{L}[q](z) - \mathcal{L}[q](w)}{z-w}$$

is the Laplace transformation of the distribution $q(x+y)$. Therefore using the famous criterion [14], we can show the claim.

By Lemma 3.5 and the assumption, the distribution $e^{-xs}k(s, t)$ in the two variables (s, t) is actually a square integrable function, and so $k(s, t)$ belongs to $L^1_{\text{loc}}[0, \infty)^2$. The rest of the proof is the same as that of Theorem 4.2. $\square$

Let $\mathbb{D}$ be the unit disc and $\zeta = \frac{z-1}{z+1}$. When $M \in \mathcal{HD}$, the function $\mathbb{D} \ni \zeta \mapsto M(z)$ is in the Hardy class $H^2(\mathbb{D})$, and so M uniquely factors in the form $M(z) = cB(z)S(z)F(z)$ with the following property [8, p.132]: c is a complex number of modulus 1. $B(z)$ is the Blaschke product

$$B(z) = \left(\frac{z-1}{z+1} \right)^k \prod_n \frac{|1-\beta_n^2|}{1-\beta_n^2} \cdot \frac{z-\beta_n}{z+\overline{\beta_n}},$$

where k is a non-negative integer and $\{\beta_n\}$ is a (finite or infinite) sequence in $\mathbb{H}_r$ satisfying

$$\sum_n \frac{\operatorname{Re} \beta_n}{1+|\beta_n|^2} < +\infty.$$

The singular component $S(z)$ is given by

$$S(z) = e^{-\sigma z} \exp \left[- \int_{-\infty}^{\infty} \frac{\lambda z + i}{\lambda + iz} d\mu(\lambda) \right],$$

where σ is a non-negative real number and μ is a finite singular positive measure. $B(z)$ and $S(z)$ are inner functions in the sense that $|B(z)| < 1$, $|S(z)| < 1$ for all $z \in \mathbb{H}_r$ and $|B(i\lambda)| = |S(i\lambda)| = 1$ for almost every λ .

The outer component $F(z)$ is given by

$$F(z) = \exp \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\lambda z + i}{\lambda + iz} \cdot \frac{\rho(\lambda) d\lambda}{1 + \lambda^2} \right],$$

where $\rho(\lambda) = \log |M(i\lambda)|^2$.

Theorem 4.5. *Let $M = cB(z)S(z)F(z) \in \mathcal{HD}$ be as above. Then $M \in \mathcal{HD}_2$ if and only if $B, S, F \in \mathcal{HD}_2$. Moreover,*

(1) $B \in \mathcal{HD}_2$ if and only if

$$\sum_n \operatorname{Re} \beta_n < +\infty.$$

(2) $S \in \mathcal{HD}_2$ if and only if $\sigma = 0$ and

$$\int_{-\infty}^{\infty} (1 + \lambda^2) d\mu(\lambda) < +\infty.$$

(3) $F \in \mathcal{HD}_2$ if and only if

$$\sup_{y \in \mathbb{R}} (\log \mathcal{P}[e^\rho](x + iy) - \mathcal{P}[\rho](x + iy)) = O(1), \quad (x \rightarrow +\infty),$$

$$\int_{-\infty}^{+\infty} (\log \mathcal{P}[e^\rho](x + iy) - \mathcal{P}[\rho](x + iy)) dy = O(x), \quad (x \rightarrow +\infty).$$

(4) Assume that there exist positive constants a , n , and C such that for all $\operatorname{Re} z \geq a$,

$$\frac{1}{|F(z)|} \leq C(1 + |z|)^n.$$

Then $F \in \mathcal{HD}_2$ if and only if there exists $x > a$ such that

$$\sup_{y \in \mathbb{R}} (\log \mathcal{P}[e^\rho](x + iy) - \mathcal{P}[\rho](x + iy)) < +\infty,$$

$$\int_{-\infty}^{+\infty} (\log \mathcal{P}[e^\rho](x + iy) - \mathcal{P}[\rho](x + iy)) dy < +\infty.$$

Proof. Let $E_1(z) = -\log |B(z)|^2$, $E_2(z) = -\log |S(z)|^2$ and

$$E(z) = \log \mathcal{P}[e^\rho](z) - \mathcal{P}[\rho](z) = \log \mathcal{P}[|F(i \cdot)|^2](z) - \log |F(z)|^2,$$

which are non-negative functions. Then we have

$$\begin{aligned} \frac{\mathcal{P}[|M(i \cdot)|^2](z)}{|M(z)|^2} - 1 &= e^{E_1(z) + E_2(z) + E_3(z)} - 1 \\ &= (e^{E_1(z)} - 1)(e^{E_2(z)} - 1)(e^{E_3(z)} - 1) \\ &+ (e^{E_1(z)} - 1)(e^{E_2(z)} - 1) + (e^{E_2(z)} - 1)(e^{E_3(z)} - 1) + (e^{E_1(z)} - 1)(e^{E_3(z)} - 1) \\ &+ (e^{E_1(z)} - 1) + (e^{E_2(z)} - 1) + (e^{E_3(z)} - 1) \\ &\geq (e^{E_1(z)} - 1) + (e^{E_2(z)} - 1) + (e^{E_3(z)} - 1) \\ &\geq E_1(z) + E_2(z) + E_3(z). \end{aligned}$$

Thus Corollary 3.6 and Theorem 4.2 implies that $M \in \mathcal{HD}_2$ if and only if $B, S, F \in \mathcal{HD}_2$. (3) and (4) also follows from the above computation with an easy inequality

$$x \leq e^x - 1 \leq \frac{e^r - 1}{r}x, \quad 0 \leq x \leq r.$$

(1) Clearly every inner function belongs to $\mathcal{HD}$. We have

$$-\log |B(z)|^2 = k \log \left| \frac{z+1}{z-1} \right|^2 + \sum_n \log \left| \frac{z+\bar{\beta}_n}{z-\beta_n} \right|^2.$$

Note that each term in the summation above is non-negative and integrable, and so

$$\int_{-\infty}^{\infty} -\log |B(x+iy)|^2 dy = k \int_{-\infty}^{\infty} \log \left| \frac{x+1+iy}{x-1+iy} \right|^2 dy + \sum_n \int_{-\infty}^{\infty} \log \left| \frac{x+iy+\bar{\beta}_n}{x+iy-\beta_n} \right|^2 dy.$$

Let $\beta_n = a_n + ib_n$ with $a_n, b_n \in \mathbb{R}$ and $a_n > 0$. Then we have

$$\int_{-\infty}^{\infty} \log \left| \frac{x+iy+\bar{\beta}_n}{x+iy-\beta_n} \right|^2 dy = \pi(x + a_n - |x - a_n|).$$

The integral $\int_{-\infty}^{\infty} -\log |B(x+iy)|^2 dy$ is finite if and only if $a_n < x$ except for finitely many n and $\sum_n a_n < \infty$. Thus if $B \in \mathcal{HD}_2$, we get $\sum_n a_n < \infty$.

Conversely, assume $\sum_n a_n < \infty$. Let $a = \sup_n a_n + 2$ and $x \geq a$. Then

$$\begin{aligned} \log \left| \frac{x+iy+\bar{\beta}_n}{x+iy-\beta_n} \right|^2 &= \log \frac{(x+a_n)^2 + (y-b_n)^2}{(x-a_n)^2 + (y-b_n)^2} \leq \frac{4xa_n}{(x-a_n)^2 + (y-b_n)^2} \\ &\leq \frac{4xa_n}{(x+1-a)^2}, \end{aligned}$$

and so

$$\frac{1}{|B(x+iy)|^2} \leq \exp \left[\frac{4x}{(x+1-a)^2} \left(k + \sum_n a_n \right) \right].$$

Thus there exists a positive constant r such that $|B(x+iy)|^{-2} \leq r$. As we have

$$\frac{1}{|B(x+iy)|^2} - 1 \leq -\frac{e^r - 1}{r} \log |B(x+iy)|^2,$$

we get $B \in \mathcal{HD}_2$.

(2) We have

$$-\log |S(x+iy)|^2 = 2\sigma x + 2 \int_{-\infty}^{\infty} \frac{x(1+\lambda^2)}{x^2 + (y-\lambda)^2} d\mu(\lambda).$$

Since

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{x(1+\lambda^2)}{x^2 + (y-\lambda)^2} d\mu(\lambda) dy = \pi \int_{-\infty}^{\infty} (1+\lambda^2) d\mu(\lambda),$$

if $S \in \mathcal{HD}_2$, then $\sigma = 0$ and $(1+\lambda^2)\mu$ is a finite measure.

Conversely assume that $\sigma = 0$ and $(1+\lambda^2)\mu$ is a finite measure. Then

$$-\log |S(x+iy)|^2 \leq \frac{2}{x} \int_{-\infty}^{\infty} (1+\lambda^2) d\mu(\lambda),$$

and there exists a positive constant l such that $|S(x+iy)|^{-2} \leq l$ for all $x \geq 1$. In the same way as above, this implies $S \in \mathcal{HD}_2$. $\square$

Corollary 4.6. *If $M_1, M_2 \in \mathcal{HD}_2$ are outer functions, then $M_1^\theta M_2^{1-\theta} \in \mathcal{HD}_2$ for all $0 \leq \theta \leq 1$.*

Proof. Hölder inequality implies

$$\begin{aligned} & \frac{\mathcal{P}[|M_1^\theta(i \cdot) M_2^{1-\theta}(i \cdot)|^2](x+iy)}{|M_1(x+iy)|^{2\theta} |M_2(x+iy)|^{2-2\theta}} - 1 \\ & \leq \left(\frac{\mathcal{P}[|M_1(i \cdot)|^2](x+iy)}{|M_1(x+iy)|^2} \right)^\theta \left(\frac{\mathcal{P}[|M_2(i \cdot)|^2](x+iy)}{|M_2(x+iy)|^2} \right)^{1-\theta} - 1 \\ & \leq \left(\frac{\mathcal{P}[|M_1(i \cdot)|^2](x+iy)}{|M_1(x+iy)|^2} \right) \left(\frac{\mathcal{P}[|M_2(i \cdot)|^2](x+iy)}{|M_2(x+iy)|^2} \right) - 1 \\ & = \left(\frac{\mathcal{P}[|M_1(i \cdot)|^2](x+iy)}{|M_1(x+iy)|^2} - 1 \right) \left(\frac{\mathcal{P}[|M_2(i \cdot)|^2](x+iy)}{|M_2(x+iy)|^2} - 1 \right) \\ & + \left(\frac{\mathcal{P}[|M_1(i \cdot)|^2](x+iy)}{|M_1(x+iy)|^2} - 1 \right) + \left(\frac{\mathcal{P}[|M_2(i \cdot)|^2](x+iy)}{|M_2(x+iy)|^2} - 1 \right). \end{aligned}$$

$\square$

We show, by example, that not every $M \in \mathcal{HD}$ is in $\mathcal{HD}_2$. For $\alpha \in \mathbb{R}$, we set $M(z) = (1+z)^\alpha$. Then $M \in \mathcal{HD}$ if and only if $-1/2 \leq \alpha < 1/2$. Corollary 3.6 shows that if $\alpha = -1/2$, the operator A_M does not generate a C_0 -semigroup at all.

Theorem 4.7. *Let $-1/2 < \alpha < 1/2$ and $M(z) = (1+z)^\alpha$. Then A_M generates a C_0 -semigroup satisfying the conditions (C1) and (C2) if and only if $\alpha = 0$.*

Proof. Thanks to Lemma 3.5, we have

$$\begin{aligned} & \int_{-\infty}^{\infty} \left[\frac{\mathcal{P}[|M(i \cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right] dy \\ & = \int_{-\infty}^{\infty} \left[\frac{1}{[(1+x)^2 + y^2]^\alpha} \frac{x}{\pi} \int_{-\infty}^{\infty} \frac{(1+s^2)^\alpha}{x^2 + (y-s)^2} ds - 1 \right] dy \\ & = x \int_0^{+\infty} \left[\frac{1}{[(1+\frac{1}{x})^2 + r^2]^\alpha \pi} \int_{-\infty}^{\infty} \frac{[\frac{1}{x^2} + (r+u)^2]^\alpha}{1+u^2} du - 1 \right] dr. \end{aligned}$$

Let $f_x(r)$ be the integrand of the right-hand side, which is a non-negative function. If $M \in \mathcal{HD}_2$, we have

$$\lim_{x \rightarrow +\infty} \frac{1}{x} \int_{-\infty}^{\infty} \left[\frac{\mathcal{P}[|M(i \cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right] dy = 0.$$

On the other hand,

$$\begin{aligned} \liminf_{x \rightarrow +\infty} \int_0^{+\infty} f_x(r) dr & \geq \int_0^{+\infty} \liminf_{x \rightarrow +\infty} f_x(r) dr \\ & = \int_0^{+\infty} \left[\frac{1}{(1+r^2)^\alpha \pi} \int_{-\infty}^{\infty} \frac{|r+u|^{2\alpha}}{1+u^2} du - 1 \right] dr. \end{aligned}$$

The integrand of the last term is a continuous non-negative function whose value at 0 is

$$\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{|u|^{2\alpha}}{1+u^2} du - 1 = \frac{1}{\cos(\alpha\pi)} - 1.$$

This is zero if and only if $\alpha = 0$. Therefore for $\alpha \neq 0$, we get $M \notin \mathcal{HD}_2$. When $\alpha = 0$, the operator A_M is nothing but A , which generates $\{S_t\}_{t \geq 0}$. $\square$

Remark 4.8. It is possible to show that $M(z) = (1+z)^\alpha \in \mathcal{HD}_b$ for $-1/2 < \alpha < 1/2$ (see Example 6.7 and Corollary 7.8).

Theorem 4.9. *Let $M(z) = \log^\alpha(a+z)$ with $\alpha > 0$ and $a \geq 1$. Then $M \in \mathcal{HD}_2$.*

Proof. Thanks to Corollary 4.4, it suffices to show

$$\int_{-\infty}^{\infty} \left[\frac{\mathcal{P}[|M(i \cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right] dy < \infty$$

for a fixed x . Since $M(z)$ is an outer function, we may and do assume that α is a natural number thanks to Corollary 4.6. To conclude the above integral is finite, it suffices to estimate $\mathcal{P}[|M(i \cdot)|^2](x+iy)$ and $|M(x+iy)|^2$ when y goes to $+\infty$. We assume that y is a sufficiently large positive number from now.

$$\begin{aligned} |M(x+iy)|^2 &= \left[\log^2 \sqrt{(a+x)^2 + y^2} + \arctan^2 \frac{y}{a+x} \right]^\alpha \\ &= \left[\left(\log y + \log \sqrt{1 + \frac{(a+x)^2}{y^2}} \right)^2 + \left(\frac{\pi}{2} - \arctan \frac{a+x}{y} \right)^2 \right]^\alpha \\ &= \left[\log^2 y + \frac{\pi^2}{4} + O\left(\frac{1}{y}\right) \right]^\alpha \\ &= \left[\log^2 y + \frac{\pi^2}{4} \right]^\alpha \left[1 + O\left(\frac{1}{y \log^2 y}\right) \right]. \end{aligned}$$

On the other hand

$$\begin{aligned} \mathcal{P}[|M(i \cdot)|^2](x+iy) &= \frac{x}{\pi} \int_{-\infty}^{\infty} \frac{\left[\log^2 \sqrt{a^2 + (y+\lambda)^2} + \arctan^2(\lambda+y) \right]^\alpha}{x^2 + \lambda^2} d\lambda \\ &\leq \frac{x}{\pi} \int_{-\infty}^{\infty} \frac{\left[\log^2 \sqrt{a^2 + (y+\lambda)^2} + \frac{\pi^2}{4} \right]^\alpha}{x^2 + \lambda^2} d\lambda \\ &= \sum_{k=0}^{\alpha} \binom{\alpha}{k} \left(\frac{\pi^2}{4} \right)^{\alpha-k} \frac{x}{\pi} \int_{-\infty}^{\infty} \frac{\log^{2k} \sqrt{a^2 + (y+\lambda)^2}}{x^2 + \lambda^2} d\lambda \\ &= \left[\log^2 y + \frac{\pi^2}{4} \right]^\alpha + \sum_{k=1}^{\alpha} \binom{\alpha}{k} \left(\frac{\pi^2}{4} \right)^{\alpha-k} \frac{x}{\pi} \int_{-\infty}^{\infty} \frac{\log^{2k} \sqrt{a^2 + (y+\lambda)^2} - \log^{2k} y}{x^2 + \lambda^2} d\lambda. \end{aligned}$$

Thus it suffices to show

$$\int_{-\infty}^{\infty} \frac{\log^n \sqrt{a^2 + (y+\lambda)^2} - \log^n y}{x^2 + \lambda^2} d\lambda = O\left(\frac{\log^{n-2} y}{y}\right), \quad (y \rightarrow +\infty),$$

for every even integer $n \geq 2$. We set

$$f_1(y) = \int_{-\infty}^{\infty} \frac{\log^n \sqrt{a^2 + (y + \lambda)^2} - \log^n |y + \lambda|}{x^2 + \lambda^2} d\lambda,$$

$$f_2(y) = \int_{-\infty}^{\infty} \frac{\log^n |y + \lambda| - \log^n y}{x^2 + \lambda^2} d\lambda,$$

and estimate f_1 and f_2 separately.

Let $g(\lambda) = \log^n \sqrt{a^2 + (\lambda)^2} - \log^n |\lambda|$. Then $g(\lambda) = O(\log^{n-1} |\lambda|/|\lambda|^2)$ as $|\lambda|$ tends to infinity and $g \in L^1(\mathbb{R})$. Fix a number r satisfying $0 < r < 1$. Then

$$\begin{aligned} |f_1(y)| &\leq \int_{-\infty}^{\infty} \frac{|g(\lambda + y)|}{x^2 + \lambda^2} d\lambda = \int_{-\infty}^{-ry} \frac{|g(\lambda + y)|}{x^2 + \lambda^2} d\lambda + \int_{-ry}^{\infty} \frac{|g(\lambda + y)|}{x^2 + \lambda^2} d\lambda \\ &\leq \frac{\|g\|_1}{x^2 + r^2 y^2} + \sup_{\lambda \geq (1-r)y} \{|g(\lambda)|\} \int_{-ry}^{\infty} \frac{1}{x^2 + \lambda^2} d\lambda \\ &= O\left(\frac{\log^{n-1} y}{y^2}\right), \quad (y \rightarrow +\infty). \end{aligned}$$

For f_2 we have

$$\begin{aligned} f_2(y) &= \int_{-\infty}^{\infty} \frac{[\log y + \log |1 + \frac{\lambda}{y}|]^n - \log^n y}{x^2 + \lambda^2} d\lambda \\ &= \frac{1}{y} \int_{-\infty}^{\infty} \frac{[\log y + \log |u|]^n - \log^n y}{\frac{x^2}{y^2} + (u - 1)^2} du \\ &= \sum_{k=1}^n \binom{n}{k} \frac{\log^{n-k} y}{y} \int_{-\infty}^{\infty} \frac{\log^k |u|}{\frac{x^2}{y^2} + (u - 1)^2} du \\ &= \frac{n \log^{n-1} y}{y} \int_{-\infty}^{\infty} \frac{\log |u|}{\frac{x^2}{y^2} + (u - 1)^2} du + O\left(\frac{\log^{n-2} y}{y}\right). \end{aligned}$$

Let $g_1(y)$ be the integral in the last term. To finish the proof, it suffices to show $g_1(y) = O(1/\log y)$. Indeed,

$$\begin{aligned}
|g_1(y)| &= \left| \int_{-1}^1 \frac{\log |u|}{\frac{x^2}{y^2} + (u-1)^2} du + \int_{|u| \geq 1} \frac{\log |u|}{\frac{x^2}{y^2} + (u-1)^2} du \right| \\
&= \left| \int_{-1}^1 \frac{\log |u|}{\frac{x^2}{y^2} + (u-1)^2} du - \int_{-1}^1 \frac{\log |u|}{\frac{x^2 u^2}{y^2} + (u-1)^2} du \right| \\
&\leq \frac{x^2}{y^2} \int_{-1}^1 \frac{(1+u)|\log |u||}{(1-u)[\frac{x^2}{y^2} + (1-u)^2]} du \\
&\leq \frac{x^2}{y^2} \int_{-1}^{1/2} \frac{(1+u)|\log |u||}{(1-u)^3} du + \frac{x^2}{y^2} \int_{1/2}^{1-1/y} \frac{(1+u)|\log |u||}{(1-u)^3} du \\
&\quad + \int_{1-1/y}^1 \frac{(1+u)|\log |u||}{(1-u)} du \\
&\leq \frac{x^2}{y^2} \int_{-1}^{1/2} \frac{(1+u)|\log |u||}{(1-u)^3} du + \frac{Cx^2(y-2)}{y^2} + \frac{C}{y} = O\left(\frac{1}{y}\right),
\end{aligned}$$

where

$$C = \sup_{1/2 < u < 1} \frac{(1+u)|\log |u||}{(1-u)}.$$

This finishes the proof. $\square$

We end this section by stating the following fact essentially proven in Lemma 4.1.

Corollary 4.10. *Let $M \in \mathcal{HD}_2$ with $e^{tA_M} = S_t + K_t$. Then*

$$\int_0^\infty e^{-2tx} d_t \|K_t\|_{H.S.}^2 = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \|\xi_{M,x+iy}\|^2 dy.$$

It is well known that the asymptotic behavior of the above quantity for large x determines that of $\|K_t\|_{H.S.}^2$ for small t . As we suspect that the latter might survive as a characteristic property of the E_0 -semigroup arising from e^{tA_M} (see Remark 5.8), it might be an interesting problem to determine the former for concrete examples. We will do it for a class of examples in next section.

5. A CLASS OF EXAMPLES

To introduce a class of functions in $\mathcal{HD}_2$, we give a heuristic argument first. In the sequel, for $\varphi, \psi \in L_{\text{loc}}^1[0, \infty)$ we denote by $\varphi * \psi$ the convolution

$$\varphi * \psi(x) = \int_0^x \varphi(s)\psi(x-s)ds = \int_0^x \varphi(x-s)\psi(s)ds.$$

Then we have $\mathcal{L}[\varphi * \psi](z) = \mathcal{L}[\varphi](z)\mathcal{L}[\psi](z)$ whenever the both sides make sense.

Let k be a function satisfying the condition (K2). We set $l(x, z)$ to be the Laplace transform of $k(x, y)$ in the second variable, that is $l(x, z) = (\kappa(x), e_z)$. Then the

condition (K2) implies

$$l(x+t, z) = e^{zt}l(x, z) - e^{zt} \int_0^t k(x, y)e^{-zy}dy + \int_0^t k(x, s)l(t-s, z)ds.$$

Assume for the moment that $k(x, y)$ is in C^1 -class. Differentiating the both sides of the above equation by t and setting $t = 0$, we get

$$\frac{\partial l}{\partial x}(x, z) = zl(x, z) + k(x, 0)(l(0, z) - 1).$$

Solving this equation, we obtain

$$e^{-zx}l(x, z) = l(0, z) + (l(0, z) - 1) \int_0^x e^{-zs}k(s, 0)ds,$$

which implies the following via the inverse Laplace transform:

$$1_{(x, \infty)}(y)k(x, y-x) = \varphi(y) + (\varphi * 1_{[0, x)}\psi)(y) - 1_{[0, x)}(y)\psi(y),$$

where $\varphi(x) = k(0, x)$ and $\psi(x) = k(x, 0)$. This is equivalent to

$$k(x, y) = \varphi(x+y) + \int_0^x \varphi(y+x-s)\psi(s)ds,$$

and $\psi = \varphi + \varphi * \psi$.

Now we start with φ in an appropriate function space and we *define* k by the above relation, which will turn out to satisfy the condition (K2). For two real numbers x and y , we denote $x \wedge y = \min\{x, y\}$.

Lemma 5.1. *Let $\varphi \in L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$. Then there exist a unique function $\psi \in L^1_{\text{loc}}[0, \infty)$ and a positive number a such that $e_a\psi \in L^1(0, \infty)$ and*

$$\mathcal{L}[\psi](z) = \frac{\mathcal{L}[\varphi](z)}{1 - \mathcal{L}[\varphi](z)}$$

for all z with $\text{Re } z \geq a$.

Proof. Take $\varepsilon > 0$ such that

$$\int_0^\varepsilon |\varphi(x)|dx < 1.$$

Then, for $a > 0$ we have

$$\|e_a\varphi\|_1 \leq \int_0^\varepsilon |\varphi(x)|dx + \int_\varepsilon^\infty |\varphi(x)|e^{-ax}dx \leq \int_0^\varepsilon |\varphi(x)|dx + \sqrt{\frac{e^{-2\varepsilon a}}{2a} \int_\varepsilon^\infty |\varphi(x)|^2dx}.$$

Thus we can choose large a so that $\|e_a\varphi\|_1 < 1$. Note that the n -fold convolution of $e_a\varphi$ is $e_a\varphi^{*n}$. Thus,

$$\sum_{n=1}^{\infty} e_a\varphi^{*n}$$

converges in $L^1(0, \infty)$ or equivalently

$$\sum_{n=1}^{\infty} \varphi^{*n}$$

converges in $L^1((0, \infty), e^{-ax}dx)$. We define ψ to be this limit. Then for $\operatorname{Re} z \geq a$ we get

$$\int_0^\infty \psi(x)e^{-zx}dx = \sum_{n=1}^\infty \mathcal{L}[\varphi](z)^n = \frac{\mathcal{L}[\varphi](z)}{1 - \mathcal{L}[\varphi](z)}.$$

□

We fix φ and ψ as above and set $\Phi(z) = \mathcal{L}[\varphi](z)$ and $\Psi(z) = \mathcal{L}[\psi](z)$. We assume that φ is a non-zero function. Since $\Phi - \Psi + \Phi\Psi = 0$ holds, we have $\varphi - \psi + \varphi * \psi = 0$.

Theorem 5.2. *Let $\varphi \in L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$ and let ψ be the function determined by Lemma 5.1. We set*

$$k(x, y) = \varphi(x + y) + \int_0^x \varphi(x + y - s)\psi(s)ds.$$

Then $k(x, y)$ satisfies the condition (K2).

Proof. Note that $k(x, y)$ makes sense as an element of $L^1_{\text{loc}}[0, \infty)^2$. Let $a \geq 1$ be a positive number such that $e_a\psi \in L^1(0, \infty)$.

First we show that $e^{-ax}k(x, y)$ is in $L^2(0, \infty)^2$. Indeed for the first term, we have

$$\begin{aligned} \int_0^\infty dx \int_0^\infty dy |e^{-ax}\varphi(x + y)|^2 &= \int_0^\infty e^{-2ax}dx \int_x^\infty dy |\varphi(y)|^2 \\ &= \int_0^\infty |\varphi(y)|^2 dy \int_0^y dx e^{-2ax} \\ &= \frac{1}{2a} \int_0^\infty (1 - e^{-2ay}) |\varphi(y)|^2 dy \\ &\leq \int_0^\infty |\varphi(y)|^2 (1 \wedge y) dy < +\infty, \end{aligned}$$

where we use the fact $(1 - e^{-by})/b \leq 1 \wedge y$ for all $y \geq 0$ and $b \geq 1$.

Using $\int_0^x |\psi(s)|ds \leq e^{ax} \|e_a \psi\|_1$, we get the following estimate for the second term:

$$\begin{aligned}
& \int_0^\infty dx \int_0^\infty dy |e^{-ax} \int_0^x \psi(s) \varphi(x+y-s) ds|^2 \\
&= \int_0^\infty e^{-2ax} dx \int_0^\infty dy \int_0^x \psi(r) \varphi(x+y-r) dr \int_0^x \overline{\psi(s) \varphi(x+y-s)} ds \\
&\leq \int_0^\infty e^{-2ax} dx \int_0^x |\psi(r)| dr \int_0^x |\psi(s)| ds \int_0^\infty |\varphi(x+y-r) \varphi(x+y-s)| dy \\
&\leq \int_0^\infty e^{-2ax} dx \int_0^x |\psi(r)| dr \int_0^x |\psi(s)| ds \int_0^\infty \frac{|\varphi(x+y-r)|^2 + |\varphi(x+y-s)|^2}{2} dy \\
&= \int_0^\infty e^{-2ax} dx \int_0^x |\psi(r)| dr \int_0^x |\psi(s)| ds \int_0^\infty |\varphi(x+y-s)|^2 dy \\
&= \|e_a \psi\|_1 \int_0^\infty e^{-ax} dx \int_0^x |\psi(x-s)| ds \int_0^\infty |\varphi(s+y)|^2 dy \\
&= \|e_a \psi\|_1 \int_0^\infty ds \int_s^\infty dx |\psi(x-s)| e^{-ax} \int_0^\infty |\varphi(s+y)|^2 dy \\
&= \|e_a \psi\|_1 \int_0^\infty e^{-as} ds \int_0^\infty dx |\psi(x)| e^{-ax} \int_0^\infty |\varphi(s+y)|^2 dy \\
&= \|e_a \psi\|_1^2 \int_0^\infty e^{-as} ds \int_0^\infty |\varphi(s+y)|^2 dy \\
&= \|e_a \psi\|_1^2 \frac{1}{a} \int_0^\infty (1 - e^{-ay}) |\varphi(y)|^2 dy \\
&\leq \|e_a \psi\|_1^2 \int_0^\infty |\varphi(y)|^2 (1 \wedge y) dy < +\infty.
\end{aligned}$$

Thus we get

$$\int_0^\infty \int_0^\infty |e^{-ax} k(x, y)|^2 dx dy \leq (1 + \|e_a \psi\|_1)^2 \int_0^\infty |\varphi(x)|^2 (1 \wedge x) dx.$$

As before, we define an $L^2(0, \infty)$ -valued function $\kappa(x)$ by $\kappa(x)(\cdot) = k(x, \cdot)$ and define $K_t \in \mathbb{B}(L^2(0, \infty))$ by

$$K_t f(x) = \begin{cases} (\kappa(t-x), f) & (x < t) \\ 0 & (t \leq x) \end{cases}$$

Then we get

$$\begin{aligned}
\|K_t\|^2 &\leq \|K_t\|_{H.S.}^2 = \int_0^t dx \int_0^\infty dy |k(x, y)|^2 \leq e^{2at} \int_0^\infty dx \int_0^\infty dy |e^{-ax} k(x, y)|^2 \\
&\leq e^{2at} (1 + \|e_a \psi\|_1)^2 \int_0^\infty |\varphi(y)|^2 (1 \wedge y) dy.
\end{aligned}$$

Thus for $\operatorname{Re} z > a$ and $f \in L^2(0, \infty)$, the Laplace transformation of $\{K_t\}_{t \geq 0}$ is well-defined and

$$\int_0^\infty e^{-zt} K_t f dt = \int_0^\infty e^{-zt} (\kappa(t), f) dt e_z.$$

Using the definition of κ , we get

$$\begin{aligned} & \int_0^\infty e^{-zt} (\kappa(t), f) dt \\ &= \int_0^\infty dt e^{-zt} \left(\int_0^\infty dy \varphi(t+y) f(y) + \int_0^\infty dy \int_0^t ds \psi(s) \varphi(y+t-s) f(y) \right) \\ &= \int_0^\infty f(y) dy \left(\int_0^\infty dt e^{-zt} \varphi(t+y) + \int_0^\infty \psi(s) ds \int_s^\infty dt e^{-zt} \varphi(y+t-s) \right) \\ &= \int_0^\infty f(y) dy \left(\int_0^\infty dt e^{-zt} \varphi(t+y) + \int_0^\infty \psi(s) e^{-zs} ds \int_0^\infty dt e^{-zt} \varphi(y+t) \right) \\ &= (1 + \Psi(z)) \int_0^\infty f(y) dy \int_0^\infty dt e^{-zt} \varphi(t+y) \\ &= (1 + \Psi(z)) (\eta_{\varphi, z}, f), \end{aligned}$$

where

$$\eta_{\varphi, z}(x) = \int_0^\infty e^{-zt} \varphi(t+x) dt.$$

As before we get

$$\mathcal{L}[\eta_{\varphi, z}](w) = \frac{\Phi(w) - \Phi(z)}{z - w}.$$

In the same way as in the proof of Theorem 4.2, we can show

$$\int_0^\infty ds \int_0^\infty dt e^{-sz_1 - tz_2} K_{s+t} e_w = \int_0^\infty ds \int_0^\infty dt e^{-sz_1 - tz_2} (K_s K_t + K_s S_t + S_s K_t) e_w,$$

which finishes the proof. $\square$

Thanks to Theorem 3.2 there exists $M \in \mathcal{HD}$ such that A_M is the generator of the semigroup $\{T_t = S_t + K_t\}_{t \geq 0}$ constructed above. The above argument shows that we have the relation

$$(1 + \Psi(z)) \eta_{\varphi, z}(x) = \xi_{M, z}(x).$$

Let $q \in L^2(0, \infty) \setminus D(A)$ such that $(1+z)\mathcal{L}[q](z) = M(z)$. Since the left-hand side is continuous in x , so is the right-hand side and Equation (3.2) implies that q is continuous. We set $x = 0$. Then the left hand side is $(1 + \Psi(z))\Phi(z) = \Psi(z)$. On the other hand the right-hand side is $q(0)/M(z) - 1$, which shows $1 + \Psi(z) = q(0)/M(z)$. Since Ψ is not a constant function, $q(0) \neq 0$. Now the equation $\Phi - \Psi + \Phi\Psi = 0$ implies $M(z) = q(0)(1 - \Phi(z))$.

Summing up the argument as above, we get

Theorem 5.3. *Let $\varphi \in L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$ and let $\{T_t\}_{t \geq 0}$ be the C_0 -semigroup constructed from φ by the above argument. Then $\{T_t\}_{t \geq 0}$ is a C_0 -semigroup satisfying the conditions (C1) and (C2). Let $A_M \in \mathcal{HD}_2$ be the generator of $\{T_t\}_{t \geq 0}$, and let $q \in L^2(0, \infty) \setminus D(A)$ be the function satisfying $(1+z)\mathcal{L}[q](z) = M(z)$. Then*

q is continuous at 0 and $M(z) = q(0)(1 - \mathcal{L}[\varphi](z))$. The relation between φ and q is given by

$$q(x) = q(0) \left(e^{-x} - \int_0^x \varphi(s) e^{s-x} ds \right),$$

$$\varphi = -\frac{q + q'}{q(0)}.$$

Remark 5.4. To construct an example of a C_0 -semigroup satisfying only (C1), the above argument still works for the Dirac function $\varphi(x) = r\delta_a(x)$ with $r \in \mathbb{C}$ and $a > 0$ if each argument is appropriately interpreted. Indeed, it is possible to show $M(z) = 1 - \Phi(z) = 1 - re^{-az}$ belongs to $\mathcal{HD}_b$ (Corollary 7.7).

Example 5.5. When $\varphi(x) = ce^{-dx}$ with $c \neq 0$ and $\operatorname{Re} d > 0$, by easy computation we have $\Phi(z) = c/(z+d)$, $\Psi(z) = c/(z+d-c)$, $\psi(x) = ce^{(c-d)x}$, and $k(x, y) = ce^{(c-d)x-dy}$.

In the rest of this section, we give an estimate of $\|K_t\|_{H.S.}$ for small t for the class of examples we have constructed in this section.

Lemma 5.6. Let $\varphi \in L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$. We set $\Phi = \mathcal{L}[\varphi]$, $M = 1 - \Phi \in \mathcal{HD}_2$, and $K_t = e^{tA_M} - S_t$. Then

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} \|\xi_{M, x+iy}\|^2 dy \sim \frac{1}{2x} \int_0^\infty (1 - e^{-2xs}) |\varphi(s)|^2 ds, \quad (x \rightarrow +\infty).$$

Proof. Since φ belongs to $L^1(0, \infty) + L^2(0, \infty)$, the quantity $\Phi(x + iy)$ converges to 0, uniformly in y , as x tends to $+\infty$. Thus Lemma 3.3 implies

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\infty}^{+\infty} \|\xi_{M, x+iy}\|^2 dy \\ &= \frac{1}{4\pi^2} \int_{-\infty}^\infty \frac{1}{|1 - \Phi(x + iy)|^2} \int_{-\infty}^\infty \frac{|\Phi(i\lambda) - \Phi(x + iy)|^2}{|x + iy - i\lambda|^2} d\lambda dy \\ &\sim \frac{1}{4\pi^2} \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{|\Phi(i\lambda) - \Phi(x + iy)|^2}{|x + iy - i\lambda|^2} d\lambda dy \\ &= \frac{1}{4\pi^2 x} \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{|\Phi(ixu + iy) - \Phi(x + iy)|^2}{1 + u^2} dy du. \end{aligned}$$

Let $f(s) = \varphi(s)(e^{-isxu} - e^{-sx})$. Then $f \in L^2(0, \infty)$ and

$$\Phi(ixu + iy) - \Phi(x + iy) = \int_0^\infty f(s) e^{-isy} ds.$$

Thus the Plancherel theorem implies that the last term above equals to

$$\begin{aligned} & \frac{1}{2\pi x} \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{|f(s)|^2}{1 + u^2} ds du \\ &= \frac{1}{2\pi x} \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{|\varphi(s)|^2 (1 + e^{-2sx} - 2e^{-sx} \cos(sxu))}{1 + u^2} ds du \\ &= \frac{1}{2x} \int_0^\infty (1 - e^{-2sx}) |\varphi(s)|^2 ds. \end{aligned}$$

□

Corollary 5.7. *Assume $\varphi \in L^2(0, \infty)$ and $M = 1 - \mathcal{L}[\varphi]$. Then,*

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} \|\xi_{M, x+iy}\|^2 dy \sim \frac{\|\varphi\|^2}{2x}, \quad (x \rightarrow +\infty),$$

$$\|K_t\|_{H.S.}^2 \sim \|\varphi\|^2 t, \quad (t \rightarrow +0).$$

Proof. The first statement immediately follows from Lemma 5.5, which together with Corollary 4.10 implies the second one via the Tauberian theorem [6, Chapter XIII.5, Theorem 3]. □

Remark 5.8. The above argument actually shows that $\varphi \in L^2(0, \infty)$ if and only if $\|K_t\|_{H.S.}^2 = O(t)$, $(t \rightarrow +0)$. On the other hand, we show in the subsequent paper [9] that the E_0 -semigroup arising from e^{tA_M} is of type I if and only if $\varphi \in L^2(0, \infty)$. In view of this fact, it is tempting to conjecture that there exists a cocycle conjugacy invariant of E_0 -semigroups that is computable from the asymptotic behavior of $\|K_t\|_{H.S.}^2$ for small t .

Now we treat $\varphi \in L_{\text{loc}}^1[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx) \setminus L^2(0, \infty)$. We set

$$m_\varphi(t) = \int_t^\infty |\varphi(x)|^2 dx,$$

which diverges as t tends to 0. Note that $m_\varphi(t)$ is integrable on every finite interval of the form $(0, a)$, $a > 0$ because the Fubini theorem implies

$$\int_0^a m_\varphi(s) ds = \int_0^\infty (u \wedge a) |\varphi(u)|^2 du = \int_0^a s |\varphi(s)|^2 ds + a \int_a^\infty |\varphi(s)|^2 ds.$$

Recall that a measurable function $L(t)$ on $(0, \infty)$ is said to be slowly varying at 0 if $L(t)$ is positive on a non-empty interval $(0, a)$ and for every $s > 0$,

$$\lim_{t \rightarrow +0} \frac{L(st)}{L(t)} = 1.$$

A measurable function $f(t)$ on $(0, \infty)$ is said to be regularly varying at 0 if there exist a real number α and a slowly varying function $L(t)$ such that $f(t) = t^\alpha L(t)$.

Lemma 5.9. *Let the notation be as in Lemma 5.6. Then*

$$\int_0^\infty e^{-xs} d_s \|K_s\|_{H.S.}^2 \sim \int_0^\infty e^{-xs} m_\varphi(s) ds, \quad (x \rightarrow +\infty).$$

In particular, if the function $t \mapsto \int_0^t m_\varphi(s) ds$ is regularly varying at 0,

$$\|K_t\|_{H.S.}^2 \sim \int_0^t m_\varphi(u) du = \int_0^t s |\varphi(s)|^2 ds + t \int_t^\infty |\varphi(s)|^2 ds, \quad (t \rightarrow +0).$$

Proof. First we claim that $tm_\varphi(t)$ converges to 0 as t tends to 0. Indeed, fix a positive number ε . Then for $0 < t \leq \varepsilon$

$$tm_\varphi(t) \leq t \int_\varepsilon^\infty |\varphi(s)|^2 ds + \int_t^\varepsilon s |\varphi(s)|^2 ds,$$

and

$$\limsup_{t \rightarrow +0} tm_\varphi(t) \leq \int_0^\varepsilon s|\varphi(s)|^2 ds,$$

which shows the claim. Thanks to Corollary 4.10, Lemma 5.5, and the above claim, the first statement follows from integration by part. The second statement follows from [6, Chapter XIII.5, Theorem 3]. $\square$

It is a routine work to show the following from Lemma 5.9 and [6, Chapter VIII.9].

Corollary 5.10. *Let the notation be as in Lemma 5.6 and let $|\varphi(x)| = x^{\alpha-1}L(x)$ where α is a non-zero constant with $0 \leq \alpha \leq 1/2$ and $L(x)$ is a slowly varying function at 0.*

(1) *When $\alpha = 0$,*

$$\|K_t\|_{H.S.}^2 \sim \int_0^t \frac{L(s)^2}{s} ds, \quad (t \rightarrow +0).$$

In particular, when $|\varphi(x)| \sim Cx^{-1}(\log x^{-1})^{-\beta}$, $(x \rightarrow +0)$ with $\beta > 1$ and a non-zero constant C ,

$$\|K_t\|_{H.S.}^2 \sim \frac{|C|^2}{(2\beta - 1)(\log \frac{1}{t})^{2\beta-1}}, \quad (t \rightarrow +0).$$

(2) *When $0 < \alpha < 1/2$,*

$$\|K_t\|_{H.S.}^2 \sim \frac{t^{2\alpha}L(t)^2}{2\alpha(1 - 2\alpha)}, \quad (t \rightarrow +0).$$

(3) *When $\alpha = 1/2$,*

$$\|K_t\|_{H.S.}^2 \sim t \int_t^\infty \frac{L(s)^2}{s} ds, \quad (t \rightarrow +0).$$

In particular, when $|\varphi(x)| \sim Cx^{-1/2}(\log x^{-1})^\beta$, $(x \rightarrow +0)$ with $\beta \geq -1/2$ and a non-zero constant C ,

$$\|K_t\|_{H.S.}^2 \sim \begin{cases} \frac{|C|^2 t \left(\log \frac{1}{t}\right)^{2\beta+1}}{2\beta+1} & (\beta > -\frac{1}{2}) \\ |C|^2 t \log \log \frac{1}{t} & (\beta = -\frac{1}{2}) \end{cases}, \quad (t \rightarrow +0).$$

6. OFF WHITE NOISE MODELS

The purpose of this section is to clarify the relationship between Tsirelson's off white noise and our semigroup $\{T_t\}_{t \geq 0}$.

Let $M \in \mathcal{HD}$ be an outer function satisfying $\overline{M(\bar{z})} = M(z)$. We introduce a measure ν on $\mathbb{R}$ by setting

$$\nu(d\lambda) = \frac{|M(i\lambda)|^2}{2\pi} d\lambda.$$

Let $G := L^2(\mathbb{R}, \nu)$. We denote by $\langle \cdot, \cdot \rangle_G$ and $\|\cdot\|_G$ the inner product and the norm of G . Let F be the unitary from $L^2(\mathbb{R})$ to G given by $Ff(\lambda) = \hat{f}(\lambda)/M(i\lambda)$, where $\hat{f}$ is the Fourier transform of f

$$\hat{f}(\lambda) = \mathcal{F}[f](\lambda) = \int_{-\infty}^{\infty} f(s)e^{-is\lambda} ds.$$

We regard $L^2(0, \infty)$ as a subspace of $L^2(\mathbb{R})$ and so $Ff(\lambda) = \mathcal{L}[f](i\lambda)/M(i\lambda)$ for $f \in L^2(0, \infty)$. We denote by U_t the bilateral shift on $L^2(\mathbb{R})$, that is, $U_t f(x) = f(x-t)$ and set $V_t = FU_t F^{-1}$. Then V_t is given by the multiplication operator of $e^{-i\lambda t}$.

Let $\mathcal{I}_0$ be the set of finite open intervals of $\mathbb{R}$ and let $\mathcal{I}$ be the set of (finite or infinite) open intervals of $\mathbb{R}$. We denote by $\mathcal{S} = \mathcal{S}(\mathbb{R})$ the set of rapidly decreasing C^∞ functions on $\mathbb{R}$. For $I \in \mathcal{I}$, we set $\mathcal{S}_I$ to be the linear span of

$$\{1_J f; f \in \mathcal{S}, J \subset I, J \in \mathcal{I}\}.$$

Lemma 6.1. *Let $I \in \mathcal{I}$ and $f \in \mathcal{S}$. Then $\mathcal{F}[1_I f] \in G$.*

Proof. Let $f \in \mathcal{S}$ and I be a finite interval (a, b) . Then

$$\mathcal{F}[1_I f](\lambda) = \frac{\mathcal{F}[1_I f'](\lambda) + f(a)e^{-ia\lambda} - f(b)e^{-ib\lambda}}{i\lambda}.$$

Since $M(z)/(1+z) \in H^2(\mathbb{H}_r)$, we conclude that $\mathcal{F}[1_I f](\lambda)M(i\lambda)$ is square integrable. The case where I is an infinite interval can be treated in the same way. $\square$

For $I \in \mathcal{I}$, we set $G_0(I) = \mathcal{F}[\mathcal{S}_I]$ and set $G(I)$ to be the closure of $G_0(I)$. By definition, we have $V_t G(I) = G(t+I)$. We denote $G_0(-\infty, \infty)$ by G_0 .

Lemma 6.2. *With the notation as above, we have $G = G(-\infty, \infty)$ and $FL^2(0, \infty) = G(0, \infty)$.*

Proof. Let $f \in G \cap G(-\infty, \infty)^\perp$. Then fM is orthogonal to $gM(i\lambda)$ in $L^2(i\mathbb{R})$ for all $g \in \mathcal{S}$. Let H_0 be the closed linear span of $\{gM\}_{g \in \mathcal{S}}$ in $L^2(i\mathbb{R})$. Then H_0 is invariant under multiplication by $e^{-it\lambda}$ for all t and so there exists a measurable subset $E \subset i\mathbb{R}$ such that $H_0 = L^2(E)$. Since $M(z)/(1+z) \in H^2(\mathbb{H}_r)$, the set of zeros of M on the imaginary axis has Lebesgue measure zero and we can conclude $H_0 = L^2(\mathbb{R})$. Thus we get $fM = 0$ and so $f = 0$. The case of the half line can be treated in a similar way by using the fact that M is an outer function and the Beurling-Lax theorem [8, p.107]. $\square$

We introduce an operator Q_0 , called the *Riesz projection*, as follows: the domain $D(Q_0)$ is G_0 and

$$Q_0 \mathcal{F}[1_I f] = \mathcal{F}[1_{I \cap (0, \infty)} f], \quad f \in \mathcal{S}, I \in \mathcal{I}.$$

Q_0 is an idempotent whose image is $G_0(0, \infty)$ and we regard it as an operator from G_0 to $G_0(0, \infty)$. When the restriction of Q_0 to $G_0(I)$ for $I \supset (0, \infty)$ is bounded, we denote by Q^I the unique bounded extension of $Q_0|_{G_0(I)}$ in $\mathbb{B}(G(I), G(0, \infty))$. We set $Q = Q^\mathbb{R}$.

Lemma 6.3. *Let $M \in \mathcal{HD}_b$ and $T_t = e^{tA_M}$. Let $f \in L^2(0, \infty)$ such that $\mathcal{L}[f](z) = \mathcal{L}[g](z)M(z)$ with $g \in \mathcal{S}_{(0, \infty)}$. Then for every $w \in \mathbb{H}_r$ in the resolvent set of A_M , we have*

$$(T_t e_w, f) = \langle F e_w, Q_0 V_{-t} F \bar{f} \rangle.$$

Proof. Easy computation yields

$$(zI - A)^{-1} e_w = \frac{e_w - e_z}{z - w},$$

and so

$$((zI - A)^{-1} e_w, f) = \frac{\mathcal{L}[g](w)M(w) - \mathcal{L}[g](z)M(z)}{z - w}.$$

Lemma 3.3 implies

$$\begin{aligned} ((zI - A_M)^{-1} e_w, f) &= (\xi_{M,z}, e_w)(e_z, f) + \frac{\mathcal{L}[g](w)M(w) - \mathcal{L}[g](z)M(z)}{z - w} \\ &= M(w) \frac{\mathcal{L}[g](w) - \mathcal{L}[g](z)}{z - w}. \end{aligned}$$

Since

$$\begin{aligned} \int_0^\infty (T_t e_w, f) e^{-tz} dt &= ((zI - A_M)^{-1} e_w, f) = M(w) \frac{\mathcal{L}[g](w) - \mathcal{L}[g](z)}{z - w} \\ &= M(w) \int_0^\infty dt \int_0^\infty g(s+t) e^{-sw-tz}, \end{aligned}$$

we get $(T_t e_w, f) = M(w) \mathcal{L}[S_t^* g](w)$.

On the other hand, using $Q_0 V_{-t} F \bar{f} = Q_0 V_{-t} \mathcal{F}[\bar{g}] = \mathcal{F}[S_t^* \bar{g}]$, we get

$$\begin{aligned} \langle F e_w, Q_0 V_{-t} F \bar{f} \rangle_G &= \frac{1}{2\pi} \int_{-\infty}^\infty \frac{1}{(w + i\lambda)M(i\lambda)} \overline{\mathcal{F}[S_t^* \bar{g}](\lambda)} |M(i\lambda)|^2 d\lambda \\ &= \frac{1}{2\pi} \int_{-\infty}^\infty \frac{M(-i\lambda) \mathcal{F}[S_t^* g](-\lambda)}{w + i\lambda} d\lambda \\ &= \frac{-1}{2\pi} \int_{-\infty}^\infty \frac{M(i\lambda) \mathcal{L}[S_t^* g](i\lambda)}{i\lambda - w} d\lambda. \end{aligned}$$

Note that $M(z)/(1+z)$ and $\mathcal{L}[S_t^* g](z)$ belong to $H^2(\mathbb{H}_r)$ and

$$\frac{-1}{2\pi} \int_{-\infty}^\infty \frac{M(i\lambda) \mathcal{L}[S_t^* g](i\lambda)}{i\lambda - w} d\lambda = \lim_{\varepsilon \rightarrow +0} \frac{-1}{2\pi i} \int_{\varepsilon - i\infty}^{\varepsilon + i\infty} \frac{M(z) \mathcal{L}[S_t^* g](z)}{z - w} dz.$$

The residue theorem implies that for $0 < \varepsilon < \operatorname{Re} w < r$ the following holds:

$$\begin{aligned} &\frac{-1}{2\pi i} \int_{\varepsilon - i\infty}^{\varepsilon + i\infty} \frac{M(z) \mathcal{L}[S_t^* g](z)}{z - w} dz \\ &= M(w) \mathcal{L}[S_t^* g](w) + \frac{-1}{2\pi i} \int_{r - i\infty}^{r + i\infty} \frac{M(z) \mathcal{L}[S_t^* g](z)}{w - z} dz. \end{aligned}$$

Since the second term of the right-hand side tends to 0 as r goes to $+\infty$, we get the statement. $\square$

Let P be the (orthogonal) projection from $G(-\infty, \infty)$ onto $G(0, \infty)$. We denote by P^I the restriction of P to $G(I)$ for $I \supset (0, \infty)$.

Theorem 6.4. *Let $M \in \mathcal{HD}$ be an outer function with $\overline{M(\bar{z})} = M(z)$.*

(1) *The following conditions are equivalent:*

(i) $M \in \mathcal{HD}_b$.

(ii) *For all $t > 0$, the restriction of Q_0 to $G_0(-t, \infty)$ is bounded.*

When these conditions hold,

$$F e^{tA_M} F^{-1} = V_t Q^{(-t, \infty)*} |_{G(0, \infty)},$$

and in particular

$$\|e^{tA_M} - S_t\| = \|Q^{(-t, \infty)} - P^{(-t, \infty)}\|.$$

(2) *The following conditions are equivalent:*

(i) $M \in \mathcal{HD}_2$.

(ii) *For all $t > 0$, the restriction of Q_0 to $G_0(-t, \infty)$ is bounded and $Q^{(-t, \infty)} - P^{(-t, \infty)}$ is a Hilbert-Schmidt operator.*

When these conditions hold,

$$\|e^{tA_M} - S_t\|_{H.S.} = \|Q^{(-t, \infty)} - P^{(-t, \infty)}\|_{H.S.}.$$

Proof. (1) Assume $M \in \mathcal{HD}_b$. The previous lemma shows that Q_0 restricted to $G_0(-t, \infty)$ is bounded for all $t > 0$ and $F e^{tA_M} F^{-1} = Q^{(-t, \infty)} V_{-t} |_{G(0, \infty)}$. Since

$$F S_t F^{-1} = V_t |_{G(0, \infty)} = P V_t P |_{G(0, \infty)},$$

we get $F S_t^* F^{-1} = P^{(-t, \infty)} V_{-t} |_{G(0, \infty)}$ and so $\|e^{tA_M} - S_t\| = \|Q^{(-t, \infty)} - P^{(-t, \infty)}\|$ holds.

Assume conversely that the restriction of Q_0 to $G_0(-t, \infty)$ is bounded for all $t > 0$ and we set $T_t = F^{-1} V_t Q^{(-t, \infty)*} F$. Since $Q^{(-t, \infty)} V_{-t} V_t |_{G_0(0, \infty)} = I$ and

$$Q^{(-t, \infty)} V_{-t} Q^{(-s, \infty)} V_{-s} |_{G_0(0, \infty)} = Q^{(-s-t, \infty)} V_{-(s+t)} |_{G_0(0, \infty)}, \quad s, t > 0,$$

$\{T_t\}_{t \geq 0}$ is a C_0 -semigroup satisfying the condition (C1). The proof of the previous lemma implies

$$\int_0^\infty (T_t e_w, f) e^{-zt} dt = ((zI - A_M)^{-1} e_w, f)$$

and so $M \in \mathcal{HD}_b$ and $T_t = e^{tA_M}$.

(2) follows from (1). □

The condition in (2) that $Q^{(-t, \infty)} - P^{(-t, \infty)}$ is a Hilbert-Schmidt operator is equivalent to that the orthogonal projection from $G(0, \infty)$ to $G(-t, 0)$ is a Hilbert-Schmidt operator. Indeed, the following statement holds in general:

Lemma 6.5. *Let H be a Hilbert space which is a topological direct sum of two closed subspaces H_1 and H_2 and let P_i be the orthogonal projection onto H_i for $i = 1, 2$. Let Q be the (not necessarily orthogonal) projection from H onto H_1 with respect to the decomposition $H = H_1 \oplus H_2$. Then the following two conditions are equivalent:*

(1) *The orthogonal projection from H_1 to H_2 is a Hilbert-Schmidt operator.*

(2) *$Q - P_1$ is a Hilbert-Schmidt operator.*

Proof. [16, p.308] implies that H has an orthogonal decomposition $H = K_1 \oplus K_2 \oplus (K_3 \oplus K_3)$ such that

$$P_1 = 1 \oplus 0 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix},$$

$$P_2 = 0 \oplus 1 \oplus \begin{bmatrix} c^2 & cs \\ cs & s^2 \end{bmatrix},$$

where $c \in \mathbb{B}(K_3)$ is a positive contraction and $s \in \mathbb{B}(K_3)$ is a positive invertible contraction with $c^2 + s^2 = I$. Then Q is given by

$$Q = 1 \oplus 0 \oplus \begin{bmatrix} 1 & -s^{-1}c \\ 0 & 0 \end{bmatrix}.$$

Thus we get $\|P_2 P_1\|_{H.S.}^2 = \text{Tr}(c^2)$ and $\|Q - P_1\|_{H.S.}^2 = \text{Tr}(s^{-2}c^2)$. $\square$

Note that the following three conditions are equivalent: (1) Q_0 is bounded. (2) The restriction of Q_0 to a Q_0 -invariant dense subspace G_1 of G_0 is bounded. (3) $G = G(-\infty, 0) \oplus G(0, \infty)$ topologically. As G_1 , we follow Tsirelson [18] and adopt the linear span of

$$\left\{ \frac{1}{(1+i\lambda)^m}, \frac{1}{(1-i\lambda)^n} \right\}_{m,n \in \mathbb{N}}.$$

In [18], B. Tsirelson observed that questions around Q_0 are reduced to those of “past-and-future geometry” via the conformal transformation

$$e^{i\theta} = \frac{i\lambda - 1}{i\lambda + 1}.$$

Let $N(e^{i\theta}) = M(i\lambda)$ and let $\mu(d\theta) = |N(e^{i\theta})|^2 d\theta$, which is a *finite* measure on the unit circle $\mathbb{T}$. Let $\tilde{G} = L^2(\mathbb{T}, \mu)$ and we set $\tilde{G}_+$ and $\tilde{G}_-$ the closed linear spans of $\{e^{in\theta}\}_{n \geq 0}$ and $\{e^{in\theta}\}_{n < 0}$ respectively. As we have

$$\mu(d\theta) = \frac{2|M(i\lambda)|^2}{1+\lambda^2} d\lambda = \frac{4\pi}{1+\lambda^2} \nu(d\lambda),$$

we can introduce a unitary W from $\tilde{G}$ onto G by setting

$$Wf(\lambda) = \frac{2\sqrt{\pi}f(\frac{1-i\lambda}{1+i\lambda})}{1+i\lambda},$$

which maps $\tilde{G}_+$ and $\tilde{G}_-$ onto $G(0, \infty)$ and $G(-\infty, 0)$ respectively. Those measures μ for which $\tilde{G}$ is a topological direct sum of $\tilde{G}_+$ and $\tilde{G}_-$ are known as Helson-Szegö measures [10, Chapter VII,D].

We get the following two corollaries from Tsirelson’s observation.

Corollary 6.6. *Let $M \in \mathcal{HD}$ be an outer function with $\overline{M(\bar{z})} = M(z)$. Then the following conditions are equivalent:*

- (1) $M \in \mathcal{HD}_b$ and $\sup_{t>0} \{ \|e^{tA_M} - S_t\| \} < \infty$.
- (2) The Riesz projection Q_0 is bounded.
- (3) The measure μ satisfies the Helson-Szegö condition: there are real functions $u, v \in L^\infty(\mathbb{T})$ with $\|v\|_\infty < \pi/2$ such that $|N(e^{i\theta})|^2 = e^{u(\theta)+\tilde{v}(\theta)}$, where $\tilde{v}$ is the conjugate function of v .

- (4) *The measure μ satisfies the Hunt-Muckenhoupt-Wheeden condition: there exists a positive constant c such that for all interval I in $\mathbb{T}$ the following holds*

$$\left[\int_I |N(e^{i\theta})|^2 d\theta \right] \left[\int_I \frac{d\theta}{|N(e^{i\theta})|^2} \right] \leq c|I|^2.$$

Example 6.7. When $M(z) = (1+z)^\alpha$ with $-1/2 < \alpha < 1/2$, we have

$$|N(e^{i\theta})|^2 = 2^{1-2\alpha} \left| \sin \frac{\theta}{2} \right|^{-2\alpha},$$

and so it is easy to show that $M(z)$ satisfies the above (4). In Corollary 7.6, we directly show (1) as well.

Corollary 6.8. *Let $M \in \mathcal{HD}$ with $\overline{M(\bar{z})} = M(z)$. Then the following conditions are equivalent:*

- (1) *M is an outer function, Q_0 is bounded and $Q-P$ is a Hilbert-Schmidt operator.*
(2) *M is an outer function satisfying*

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{|\log |M(is)| - \log |M(it)||^2}{|s-t|^2} ds dt < \infty.$$

(3)

$$\sup_{x>0} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{|M(x+iy) - M(is)|^2}{[x^2 + (y-s)^2] |M(x+iy)|^2} ds dy < \infty.$$

(4)

$$\sup_{x>0} \frac{1}{x} \int_{-\infty}^{\infty} \left(\frac{\mathcal{P}[|M(i\cdot)|^2](x+iy)}{|M(x+iy)|^2} - 1 \right) dy < \infty.$$

- (5) *$M \in \mathcal{HD}_2$ and $\sup_{t>0} \{ \|e^{tA_M} - S_t\|_{H.S.} \} < \infty$.*

Proof. Tsirelson [18, Theorem 3.2] (with Lemma 6.5) shows that (2) implies (1). The equivalence of (3), (4), and (5) has already been shown. Theorem 6.4 shows that (1) implies (5).

Assume (1). Since μ is a finite measure, Ibragimov-Solev theorem (see [18, Proposition 1.8]) implies that $\log |N(e^{i\theta})|^2$ belongs to the Sobolev space $W_2^{1/2}(\mathbb{T})$ and equivalently, the function M satisfies the condition of (2).

Assume (3), (4) and (5) now. (4) implies that $M(z)$ has no zeros on $\mathbb{H}_r$ and the Blaschke component of $M(z)$ is trivial. (3) with the argument of the proof of Theorem 4.5 shows that M has a trivial singular inner component as well and so $M(z)$ is outer. Corollary 6.6 implies that the Riesz projection is bounded and Q is well-defined. Let $P_{-(t,\infty)}$ be the orthogonal projection from G onto $G(-t, \infty)$. Note that $P_{-(t,\infty)}$ converges to I in the strong operator topology as t tends to $+\infty$. (5) implies that there is a positive constant c such that $\|(Q-P)P_{-(t,\infty)}\|_{H.S.} \leq c$ for all $t > 0$. Since the trace of $\mathbb{B}(G)$ is lower semi-continuous in the weak operator topology, we get

$$\|Q-P\|_{H.S.}^2 = \text{Tr}((Q-P)(Q-P)^*) = \lim_{t \rightarrow +\infty} \text{Tr}((Q-P)P_{-(t,\infty)}(Q-P)^*) \leq c.$$

Thus $Q-P$ is a Hilbert-Schmidt operator and (1) holds. $\square$

We have seen that $M(z) = \log^\alpha(a + z)$ belongs to $\mathcal{HD}_2$ for $a \geq 1$ and $\alpha > 0$ in Section 4. Using Tsirelson's criterion [18, Proposition 3.6], we can actually show that M satisfies the conditions of Corollary 6.7 if and only if $a > 1$ (α can be an arbitrary real number). $\log^\alpha(1 + z)$ with $\alpha > 0$ is a typical example of $M \in \mathcal{HD}_2$ without satisfying the conditions of Corollary 6.7. This is caused by a zero of M on the imaginary axis. Indeed, for every $M \in \mathcal{HD}_2$, the function $M(z)z/(1 + z)$ belongs to $\mathcal{HD}_2$ and does not satisfy the conditions of Corollary 6.7 (See Proposition 7.11.)

We end this section with showing that the Tsirelson's E_0 -semigroups constructed from off white noises in [17] actually come from our construction in Theorem 2.3. Although it is possible to show the statement by a purely measure theoretical argument as in [17] (in fact, the author first obtained the statement in that way), we take an operator theoretical approach using Shale's result [15] inspired by Bhat and Srinivasan's paper [4].

Let $\rho(x)$ be a real valued measurable function on $\mathbb{R}$ such that

$$\int_{-\infty}^{+\infty} \frac{e^{\rho(\lambda)}}{1 + \lambda^2} < \infty,$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{|\rho(s) - \rho(t)|^2}{|s - t|^2} ds dt < \infty.$$

We set

$$M(z) = \exp \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\lambda z + i}{\lambda + iz} \cdot \frac{\rho(\lambda) d\lambda}{1 + \lambda^2} \right],$$

and set $\{T_t\}_{t \geq 0}$ to be the C_0 -semigroup corresponding to M . In the above setting, we have

$$\nu(d\lambda) = \frac{e^{\rho(\lambda)}}{2\pi} d\lambda.$$

Thus our ν is nothing but a multiple of ν in [17, Section 9] and $G(s, t)$ is the complexification of $G_{s,t}$ in [17, Section 9]. We set $G(s, t)_{\mathbb{R}}$ to be the real part of $G(s, t)$, that is, $G(s, t)_{\mathbb{R}}$ is the closure of the linear span of the functions of the form $\mathcal{F}[1_I f]$ where $I \subset (s, t)$ and $f \in \mathcal{S}$ is real valued.

Let

$$L : G(-\infty, 0) \oplus G(0, \infty) \ni f \oplus g \mapsto f + g \in G,$$

where we regard $G(-\infty, 0) \oplus G(0, \infty)$ as an orthogonal direct sum. Since $I - L^*L$ is a Hilbert-Schmidt class operator, Shale's result [15] shows that there exists a unitary operator $\Gamma(L) : e^{G(-\infty, 0)} \otimes e^{G(0, \infty)} \rightarrow e^G$ satisfying

$$\begin{aligned} \Gamma(L)(W(f_1 + ig_1) \otimes W(f_2 + ig_2))\Gamma(L)^* &= W(L(f_1 \oplus f_2) + iL^{*-1}(g_1 \oplus g_2)) \\ &= W(f_1 + f_2 + iL^{*-1}(g_1 \oplus g_2)), \end{aligned}$$

where $f_1, g_1 \in G(-\infty, 0)_{\mathbb{R}}$ and $f_2, g_2 \in G(0, \infty)_{\mathbb{R}}$. Then Tsirelson's E_0 -semigroup β_t acting on $\mathbb{B}(e^{G(0, \infty)})$ is given by

$$1 \otimes \beta_t(W(f + ig)) = \text{Ad}(\Gamma(L)\Gamma(V_t)\Gamma(L)^*)(W(f + ig)).$$

Theorem 6.9. *Let the notation be as above. Then Tsirelson's E_0 -semigroup $\{\beta_t\}_{t \geq 0}$ is conjugate to the E_0 -semigroup $\{\alpha_t\}_{t \geq 0}$ acting on $\mathbb{B}(e^{L^2(0, \infty)})$ given by*

$$\alpha_t(W(f + ig)) = W(S_t f + iT_t g), \quad f, g \in L^2(0, \infty)_{\mathbb{R}}.$$

Proof. Direct computation shows that for $f, g \in G(0, \infty)_{\mathbb{R}}$, we have

$$\begin{aligned} 1 \otimes \beta_t(W(f + ig)) &= W(L^{-1}V_t f + iL^*V_t L^{*-1}(0 \oplus g)) \\ &= W((0 \oplus V_t f) + iL^*V_t L^{*-1}(0 \oplus g)). \end{aligned}$$

Let $f_1 \in G(-\infty, 0)_{\mathbb{R}}$ and $f_2 \in G(0, \infty)_{\mathbb{R}}$. Then

$$\begin{aligned} \langle L^*V_t L^{*-1}(0 \oplus g), f_1 \oplus f_2 \rangle &= \langle 0 \oplus g, L^{-1}V_{-t}L(f_1 \oplus f_2) \rangle \\ &= \langle 0 \oplus g, L^{-1}V_{-t}(f_1 + f_2) \rangle \\ &= \langle 0 \oplus g, V_{-t}f_1 + (I - Q^{(-t, \infty)})V_{-t}f_2 \oplus Q^{(-t, \infty)}V_{-t}f_2 \rangle \\ &= \langle g, Q^{(-t, \infty)}V_{-t}f_2 \rangle. \end{aligned}$$

Thus Lemma 6.3 shows the statement. $\square$

Remark 6.10. Let $\varphi \in L^1_{\text{loc}}[0, \infty) \cap L^2((0, \infty), (1 \wedge x)dx)$ with $\|\varphi\|_1 < 1$ and we set $M(z) = 1 - \mathcal{L}[\varphi](z)$. Then there exist positive constants $0 < c_1 \leq 1 \leq c_2$ such that $c_1 \leq |M(i\lambda)| \leq c_2$ holds for all λ . As a system of topological vector spaces, $\{G(s, t)\}_{s < t}$ and $\{L^2(s, t)\}_{s < t}$ are canonically isomorphic. Since Tsirelson's infinitesimal sequence invariant in [17] is rather an invariant for $\{G(s, t)\}_{s < t}$ as a system of topological vector spaces, it does not distinguish the E_0 -semigroup arising from M from the CCR flow. Yet, we will show that even such M sometimes produces a E_0 -semigroup of type III in the forthcoming paper [9]. Namely we will show that the resulting E_0 -semigroup is of type III if and only if $\varphi \notin L^2(0, \infty)$ and that there are uncountably many mutually non cocycle conjugate E_0 -semigroups of type III arising in this way.

7. APPENDIX.

In this appendix, we show how to compute $K_t = e^{tA_M} - S_t$ for a function M in $\mathcal{HD}_b$ that is not necessarily in $\mathcal{HD}_2$ and present computation of concrete examples.

Lemma 7.1. *Let $M \in \mathcal{HD}$ and $f \in D(A)$. Then*

$$(\xi_{M,z}, f) = \frac{1}{M(z)} \int_0^\infty e^{-tz} (q, S_t(f' - f)) dt,$$

where $q \in L^2(0, \infty) \setminus D(A)$ with $M(z) = (1 + z)\mathcal{L}[q](z)$.

Proof. Thanks to Equation (3.2), we get

$$\begin{aligned} M(z)(\xi_{M,z}, f) &= (q, f) - (1 + z) \int_0^\infty e^{-tz} (q, S_t f) dt \\ &= (q, f) - (1 + z)(q, (zI - A)^{-1}f) = -(q, (zI - A)^{-1}(A + I)f) \\ &= \int_0^\infty e^{-tz} (q, S_t(f' - f)) dt. \end{aligned}$$

$\square$

Lemma 7.2. *Let $M \in \mathcal{HD}$ such that there exist a natural number n and positive constants a and C such that $|M(z)| \geq C(1 + |z|)^{-n}$ for $\operatorname{Re} z > a$.*

- (1) *There exists a distribution $k \in \mathcal{D}'(\mathbb{R} \times (0, \infty))$ with support in $[0, \infty) \times (0, \infty)$ such that for every $f \in \mathcal{D}(0, \infty)$, $g \in \mathcal{D}(\mathbb{R})$ and $b > a$,*

$$\frac{1}{2\pi i} \int_{b-i\infty}^{b+i\infty} \left(\int_{-\infty}^{\infty} e^{xz} g(x) dx (\xi_{M,z}, f) \right) dz = (k, g \otimes f).$$

- (2) *Moreover, if there exists $r \in L^2(0, \infty)$ such that $\mathcal{L}[r](z) = 1/[(1+z)M(z)]$, then $(k, g \otimes f) = (k_f, g)$ where*

$$k_f(x) = -(q, f - f')r(x) - \int_0^x r(s)(q, S_{x-s}(f - 2f' + f''))ds.$$

Proof. (1) Let $l(t) = (q, S_t(f' - f))$, which is a bounded smooth function on $[0, \infty)$ vanishing at infinity. Since there exists positive constants a and $C' > 0$ such that

$$|(\xi_{M,z}, f)| \leq C'(1 + |z|)^n, \quad \operatorname{Re} z > a,$$

there exists a distribution $k_f \in \mathcal{D}'(\mathbb{R})$ with support in $[0, \infty)$ such that the integral in the statement (1) is (k_f, g) . Note that it does not depend on $b > a$. By definition, for fixed f the map $\mathcal{D}(\mathbb{R}) \ni g \mapsto (k_f, g)$ is continuous. We claim that for fixed g , the map $\mathcal{D}(0, \infty) \ni f \mapsto (k_f, g)$ is continuous. Note that the function

$$y \mapsto \int_{-\infty}^{\infty} e^{x(b+iy)} g(x) dx,$$

is rapidly decreasing. Since

$$|(\xi_{M,b+iy}, f)| \leq C'(1 + |b + iy|)^n \|q\| (\|f\| + \|f'\|),$$

there exists a constant C_g , depending on g , such that $|(k_f, g)| \leq C_g(\|f\| + \|f'\|)$ holds for all $f \in \mathcal{D}(0, \infty)$, which shows the claim. Thus the Schwartz nuclear theorem [14] implies the statement.

- (2) Since

$$(\xi_{M,z}, f) = (1+z)\mathcal{L}[r](z)\mathcal{L}[l](z) = \mathcal{L}[r](z)(\mathcal{L}[l+l'](z) + l(0)),$$

we get $k_f = r * (l + l') + (q, f' - f)r$. □

Corollary 7.3. *Let M , k , q , and r be as above.*

- (1) *If the derivatives q' and q'' in $\mathcal{D}'(0, \infty)$ belong to $L^2_{\text{loc}}(0, \infty)$, then k is in $L^1_{\text{loc}}(0, \infty)^2$ and is given by*

$$\begin{aligned} k(x, y) &= -r(x)(q(y) + q'(y)) \\ &\quad - \int_0^x r(s)(q(x+y-s) + 2q'(x+y-s) + q''(x+y-s))ds. \end{aligned}$$

- (2) *If r' and q' in $\mathcal{D}'(0, \infty)$ belong to $L^2_{\text{loc}}(0, \infty)$ and r is continuous at 0, then k is in $L^1_{\text{loc}}(0, \infty)^2$ and is given by*

$$\begin{aligned} k(x, y) &= -r(0)(q(x+y) + q'(x+y)) \\ &\quad - \int_0^x (r(s) + r'(s))(q(x+y-s) + q'(x+y-s))ds. \end{aligned}$$

Theorem 7.4. *Let $M \in \mathcal{HD}$ with positive constants n , a , and C as in Lemma 7.2 and let k be the distribution defined in Lemma 7.2. Then $M \in \mathcal{HD}_b$ if and only if there exist positive constants D and b such that for all $t \geq 0$,*

$$|(k, g \otimes f)| \leq De^{bt} \|f\|_2 \cdot \|1_{(0,\infty)} g\|_2, \quad f \in \mathcal{D}(0, \infty), \quad g \in \mathcal{D}((-\infty, t)).$$

When this assumption is satisfied, the semigroup $\{e^{tA_M} = S_t + K_t\}_{t \geq 0}$ is given as follows: Let $\tilde{K}_t \in \mathbb{B}(L^2(0, \infty), L^2(0, t))$ determined by $(k, g \otimes f) = (\tilde{K}_t f, 1_{(0,t)} g)$. Then

$$K_t f(x) = \begin{cases} \tilde{K}_t f(t-x) & (x < t) \\ 0 & (t \leq x) \end{cases}$$

Proof. When $M \in \mathcal{HD}_b$, necessity of the constants D and b with the above property is obvious.

Assume conversely that there exist $D > 0$ and b satisfying the conditions above and define $\tilde{K}_t$ and K_t as above. The family $\{K_t\}_{t \geq 0}$ is strongly continuous with $\|K_t\| \leq De^{bt}$ and the Laplace transformation of $\{K_t\}_{t \geq 0}$ exists for $\operatorname{Re} z > b$. Let $f \in \mathcal{D}(0, \infty)$ and $g \in \mathcal{D}(\mathbb{R})$. Note that for $0 < s < t$, $\tilde{K}_s f(x) = \tilde{K}_t f(x)$ for almost all $x \in (0, t)$. We define $Kf(x)$ to be $\tilde{K}_t f(x)$ with $x < t$. Note that we have

$$\int_0^t |Kf(x)|^2 dx \leq D^2 e^{2bt} \|f\|^2.$$

Then

$$\begin{aligned} \int_0^\infty e^{-tz} (K_t f, g) dt &= \int_0^\infty dt e^{-tz} \int_0^t Kf(t-x) g(x) dx \\ &= \int_0^\infty dx g(x) \int_0^\infty e^{-(t+x)z} Kf(t) dt \\ &= (g, e_z) \int_0^\infty e^{-tz} Kf(t) dt \\ &= (g, e_z) \lim_{m \rightarrow \infty} (k, h_m e_z \otimes f), \end{aligned}$$

where $\{h_m\}_{m=0}^\infty$ is an increasing sequence in $\mathcal{D}(-\infty, \infty)$ converging pointwisely to a smooth function h satisfying the following properties: $0 \leq h_m(x) \leq 1$ for all $x \in \mathbb{R}$ and $h_m(x) = 1$ for $[-1, m]$ and $h_m(x) = 0$ for $x \leq -2$. We show

$$\lim_{m \rightarrow \infty} (k, h_m e_z \otimes f) = (\xi_{M,z}, f).$$

Indeed, let k_f be the distribution whose Laplace transformation is $(\xi_{M,z}, f)$. Note that $e_a k_f$ is a tempered distribution. Thus for $\operatorname{Re} z > a, b$, we have

$$\begin{aligned} \lim_{m \rightarrow \infty} (k, h_m e_z \otimes f) &= \lim_{m \rightarrow \infty} (k_f, e_z h_m) = \lim_{m \rightarrow \infty} (e_a k_f, e_{z-a} h_m) = (e_a k_f, e_{z-a} h) \\ &= \mathcal{L}[e_a k_f](z-a) = \mathcal{L}[k_f](z) = (\xi_{M,z}, f). \end{aligned}$$

Let $T_t = S_t + K_t$, whose Laplace transformation is the resolvent of A_M . Since there exists a constant $D_1 > 0$ such that $\|T_t\| \leq D_1 e^{bt}$, we have

$$\|(zI - A_M)^{-m}\| \leq \frac{D_1}{(\operatorname{Re} z - b)^m}, \quad m \in \mathbb{N}, \quad \operatorname{Re} z > b,$$

which shows that A_M generates a C_0 -semigroup thanks to [19], which should coincides with $\{T_t\}_{t \geq 0}$. $\square$

When $M(z) = (1+z)^\alpha$ with $-1/2 < \alpha < 1/2$, we have $q(x) = x^{-\alpha}e^{-x}/\Gamma(1-\alpha)$ and $r(x) = x^\alpha e^{-x}/\Gamma(1+\alpha)$ and so

$$q(x) + q'(x) = \frac{-\alpha x^{-(1+\alpha)}e^{-x}}{\Gamma(1-\alpha)},$$

$$q(x) + 2q'(x) + q''(x) = \frac{\alpha(1+\alpha)x^{-(2+\alpha)}e^{-x}}{\Gamma(1-\alpha)}.$$

Thus

$$\begin{aligned} k(x, y) &= \frac{\alpha e^{-(x+y)}}{\Gamma(1+\alpha)\Gamma(1-\alpha)} \left[x^\alpha y^{-(\alpha+1)} - (\alpha+1) \int_0^x s^\alpha (x+y-s)^{-(\alpha+2)} ds \right] \\ &= \frac{x^\alpha e^{-(x+y)}}{\Gamma(\alpha)\Gamma(1-\alpha)} \left[y^{-(\alpha+1)} - (\alpha+1)x \int_0^1 t^\alpha (x+y-xt)^{-(\alpha+2)} dt \right] \\ &= \frac{\sin(\pi\alpha)x^\alpha e^{-(x+y)}}{\pi} \left[y^{-(\alpha+1)} - (\alpha+1)x \int_1^\infty [(x+y)u-x]^{-(\alpha+2)} du \right] \\ &= \frac{\sin(\pi\alpha)x^\alpha e^{-(x+y)}}{\pi} \left[y^{-(\alpha+1)} - \frac{xy^{-(\alpha+1)}}{x+y} \right] \\ &= \frac{\sin(\pi\alpha)}{\pi} \cdot \frac{x^\alpha y^{-\alpha} e^{-(x+y)}}{x+y}. \end{aligned}$$

Lemma 7.5. *Let*

$$k(x, y) = \frac{\sin(\pi\alpha)}{\pi} \cdot \frac{x^\alpha y^{-\alpha} e^{-(x+y)}}{x+y}.$$

For $-1/2 < \alpha < 1/2$, the operator K given by

$$Kf(x) = \int_0^\infty k(x, y)f(y)dy$$

is a bounded operator in $\mathbb{B}(L^2(0, \infty))$ with $\|K\| \leq |\tan(\alpha\pi)|$.

Proof. We apply the Schur test [7, Theorem 5.2] to k . Indeed, using the fact

$$\int_0^\infty \frac{1}{(1+t)t^\beta} dt = \frac{\pi}{\sin(\beta\pi)}, \quad 0 < \beta < 1,$$

we get

$$\begin{aligned} \int_0^\infty |k(x, y)| \frac{1}{\sqrt{x}} dx &\leq \frac{|\sin(\alpha\pi)|}{\pi\sqrt{y}} \int_0^\infty \frac{dt}{(1+t)t^{1/2-\alpha}} = \frac{|\tan(\alpha\pi)|}{\sqrt{y}}, \\ \int_0^\infty |k(x, y)| \frac{1}{\sqrt{y}} dy &\leq \frac{|\sin(\alpha\pi)|}{\pi\sqrt{x}} \int_0^\infty \frac{dt}{(1+t)t^{1/2+\alpha}} = \frac{|\tan(\alpha\pi)|}{\sqrt{x}}, \end{aligned}$$

which shows the statement. $\square$

Corollary 7.6. *Let $M(z) = (1+z)^\alpha$ with $-1/2 < \alpha < 1/2$. Then $M \in \mathcal{HD}_b$ with*

$$e^{tA_M} f(x) = 1_{(0,t)}(x) K f(t-x) + 1_{[t,\infty)}(x) f(x-t),$$

$$K f(x) = \frac{\sin(\alpha\pi)}{\pi} \int_0^\infty \frac{x^\alpha y^{-\alpha} e^{-(x+y)}}{x+y} f(y) dy.$$

Corollary 7.7. *$M(z) = 1 - re^{-az}$ with $a > 0$ and $r \in \mathbb{C}$. Then $M \in \mathcal{HD}_b$.*

Proof. Easy computation yields

$$(k, g \otimes f) = \sum_{n=1}^{\infty} r^n \int_{(n-1)a}^{na} g(x) f(na-x) dx,$$

which satisfies the assumption of Theorem 7.4. □

Proposition 7.8. *Let $M \in \mathcal{HD}_b$ such that $1/(zM(z))$ is the Laplace transformation of a function in $L^2_{\text{loc}}(0, \infty)$, that is, there exists $a > 0$ satisfying*

$$\sup_{x \geq a} \int_{-\infty}^{\infty} \frac{dy}{|x+iy|^2 |M(x+iy)|^2} < \infty.$$

Let $\beta \in \mathbb{C}$ and $\gamma \in \mathbb{H}_r$ with $\text{Re } \beta \geq 0$. Then

$$M_1(z) = \frac{z-\beta}{z+\gamma} M(z)$$

belongs to $\mathcal{HD}_b$. Let k and k_1 be the distributions defined in Lemma 7.2 for M and M_1 respectively. Then

$$k_1(x, y) = k(x, y) + (\beta + \gamma) r_\beta(x) q_\gamma(y),$$

where $r_\beta \in L^2_{\text{loc}}(0, \infty)$ and $q_\gamma \in L^2(0, \infty)$ are determined by

$$\mathcal{L}[q_\gamma](z) = \frac{M(z)}{z+\gamma}, \quad \mathcal{L}[r_\beta](z) = \frac{1}{(z-\beta)M(z)}.$$

In particular, $e^{tA_{M_1}} - e^{tA_M}$ is a rank one operator.

Proof. Thanks to Lemma 3.3, we have

$$\mathcal{L}[\xi_{M_1, z}](w) = \frac{1 - \frac{M_1(w)}{M_1(z)}}{z-w} = \mathcal{L}[\xi_{M, z}](w) + \frac{\beta + \gamma}{(z-\beta)M(z)} \cdot \frac{M(w)}{w+\gamma},$$

and so

$$\xi_{M_1, z} = \xi_{M, z} + \frac{\beta + \gamma}{(z-\beta)M(z)} q_\gamma.$$

Now the statement follows from Theorem 7.4. □

REFERENCES

- [1] Araki, H. *A lattice of von Neumann algebras associated with the quantum theory of a free Bose field*. J. Math. Phys. **4** (1963), 1343–1362.
- [2] Araki, H. *On quasifree states of the canonical commutation relations. II*. Publ. Res. Inst. Math. Sci. **7** (1971/72), 121–152.
- [3] Arveson, W. *Noncommutative Dynamics and E-semigroups*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [4] Bhat, R. Srinivasan, R. *On Product Systems arising from Sum Systems*. preprint.
- [5] van Daele, A. *Quasi-equivalence of quasi-free states on the Weyl algebra*. Comm. Math. Phys. **21** (1971), 171–191.
- [6] Feller, William, *An Introduction to Probability Theory and its Applications. Vol. II*. Second edition, John Wiley & Sons, Inc., New York-London-Sydney 1971.
- [7] Halmos, P.; Sunder, V. S. *Bounded Integral Operators on L^2 Spaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete, 96. Springer-Verlag, Berlin-New York, 1978.
- [8] Hoffman, K. *Banach Spaces of Analytic Functions*. Prentice-Hall, Inc., Englewood Cliffs, N. J. 1962.
- [9] Izumi, M. Srinivasan, R. in preparation.
- [10] Koosis, P. *Introduction to H^p Spaces. With an Appendix on Wolff's Proof of the Corona Theorem*. London Mathematical Society Lecture Note Series, 40. Cambridge University Press, Cambridge-New York, 1980.
- [11] Powers, R. T. *A nonspatial continuous semigroup of $*$ -endomorphisms of $\mathfrak{B}(\mathfrak{h})$* . Publ. Res. Inst. Math. Sci. **23** (1987), 1053–1069.
- [12] Powers, R. T. *Construction of E_0 -semigroups of $\mathfrak{B}(\mathfrak{h})$ from CP-flows*. Advances in Quantum Dynamics, 57–97, Contemp. Math., 335, Amer. Math. Soc., Providence, RI, 2003.
- [13] Price, G. (editor), *Advances in Quantum Dynamics*. Contemp. Math., 335, Amer. Math. Soc., Providence, RI, 2003.
- [14] Schwartz, Laurent. *Théorie des Distributions*. Hermann, Paris 1966.
- [15] Shale, D. *Linear symmetries of free boson fields*. Trans. Amer. Math. Soc. **103** (1962), 149–167.
- [16] Takesaki, M. *Theory of Operator Algebras. I*. Encyclopaedia of Mathematical Sciences, 124. Operator Algebras and Non-commutative Geometry, 5. Springer-Verlag, Berlin, 2002.
- [17] Tsirelson, B. *Non-isomorphic product systems*. Advances in Quantum Dynamics, 273–328, Contemp. Math., 335, Amer. Math. Soc., Providence, RI, 2003.
- [18] Tsirelson, B. *Spectral densities describing off-white noises*. Ann. Inst. H. Poincaré Probab. Statist. **38** (2002), 1059–1069.
- [19] Yosida, K. *Functional Analysis*. Sixth edition. Springer-Verlag, Berlin-New York, 1980.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY,
SAKYO-KU, KYOTO 606-8502, JAPAN

E-mail address: izumi@math.kyoto-u.ac.jp